

skin, eyes or clothing. Avoid breathing vapors. Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly with soap and water after using. Food utensils such as tablespoons should not be used for food purposes after use with insecticides. This product is a cholinesterase inhibitor. If swallowed give two tablespoons of salt in a glass of warm water to induce vomiting. Do not treat nursing animals or animals under 1 month of age. DO NOT USE WITHIN TWO WEEKS ON ANIMALS BEING TREATED WITH OTHER INSECTICIDES INCLUDING FLEA COLLARS OR INTERNAL ORGANOPHOSPHATE WORMING DRUGS. CALL A PHYSICIAN if poisoning symptoms are present. To physician or veterinarian the antidote is atropine.

ACTIVE INGREDIENTS 2-(1-Methylethoxy)-phenol methylcarbamate 8.00%
INERT INGREDIENTS 92% Catalog No. V-33

MANUFACTURED BY
FLORIDA VETERINARY LABORATORIES
DIVISION OF PET CHEMICALS, INC.
P. O. BOX 660656
MIAMI SPRINGS, FLORIDA 33166

Copyright 1975 Pat. Pend. 11775-50M EPA EST. NO. 7225-FL-1

CONCENTRATE for Dogs and Cats

Add ¼ ounce (1 tbsp.) to a quart of water

Net Contents 8 Fl. Oz.

EPA Reg. No. 4758-118-AA



CAUTION: Keep out of reach of children.
See back panel for additional cautions.

4758-118

VIP

SOLD TO VETERINARIANS ONLY

VIP DIP concentrate is almost odorless and makes a clear solution when mixed with water. VIP DIP usually kills fleas in ten minutes or less and ticks in 30 to 60 minutes.

FOR BEST FLEA AND TICK CONTROL and to help prevent re-infestation the premises should be treated. Use VIP INSECT FOGGER for the house or building and a spray or powder recommended by the Veterinarian for outside areas.

TO KILL FLEAS ON CATS, & FLEAS & TICKS ON DOGS & PUPPIES: Mix ½ oz. (one tablespoon) to a quart of water or 2 oz. (4 tablespoons) to a gallon of water. USE WARM TAP WATER. MEASURE ACCURATELY. Mix well. Sponge or sprinkle on, or dip animal until coat is thoroughly wet. Rub well into fur, between toes and in ears. For better lasting effects do not rinse but, allow to dry naturally, then comb and brush if desired. Use once a week if needed. For faster kill of engorged female ticks on dogs, 1 oz. (2 tablespoons) to a quart of water or 4 oz. (8 tablespoons) to a gallon of water may be used.

DIP

FLEA AND TICK CONCENTRATE for Dogs and Cats

Add ½ ounce (1 tbsp.) to a quart of water.

CAUTION: Keep out of reach of children.
See back panel for additional cautions.



NET CONTENTS 1 Gallon EPA Reg. No. 4758-118-AA

DO NOT RE-USE CONTAINER.

Rinse with water and pierce to destroy when empty.

CAUTION: DO NOT USE FULL STRENGTH. MIX AS DIRECTED. Keep out of reach of children. Contact with skin may cause toxic symptoms. Avoid contact with skin, eyes, or clothing. Avoid breathing vapors. Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly with soap and water after using. Food utensils such as tablespoons should not be used for food purposes after use with insecticides. This product is a cholinesterase inhibitor. If swallowed give two tablespoons of salt in a glass of warm water to induce vomiting. Do not treat nursing animals or animals under 1 month of age. DO NOT USE WITHIN TWO WEEKS ON ANIMALS BEING TREATED WITH OTHER INSECTICIDES INCLUDING FLEA COLLARS OR INTERNAL ORGANOPHOSPHATE WORMING DRUGS. CALL A PHYSICIAN if poisoning symptoms are present. To physician or veterinarian the antidote is atropine.

ACTIVE INGREDIENTS: 2-(1-Methylethoxy)-phenol methylcarbamate 8.08%

INERT INGREDIENTS: 92%

Catalog No. V34

Copyright 1975 Pat. Pend. 11775-50M EPA EST. NO. 7225-FL-1

Manufactured by
FLORIDA VETERINARY LABORATORIES
Division of Pet Chemicals, Inc.
P.O. Box 660656
Miami Springs, Florida 33166