

RESTRICTED USE PESTICIDE

Due to very high acute toxicity to Humans and Birds

For retail sale only to certified commercial aerial applicators or persons under their direct supervision. For use only by certified aerial commercial applicators. Direct supervision for this product is defined as the certified commercial aerial applicator being physically present during mixing, loading, repair and cleaning of application equipment. Certified commercial aerial applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

SUPPLEMENTAL LABELING TO

PARATHION 4 EC

EPA Reg. No. 4787-15

FOR USE ON CANOLA / RAPE (OILSEED VARIETIES ONLY)

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

THIS LABEL MUST BE IN POSSESSION OF USER AT THE TIME OF APPLICATION.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CANOLA/RAPE (OILSEED VARIETIES ONLY): For control of Cabbage Seed Pod Weevils, Aphids, Diamondback Moth Larvae, and Flea Beetles, apply 1 pint of PARATHION 4E per acre per application, in sufficient water for good coverage (3 to 10 gallons of water per acre). Apply when damaging populations are present.

- o For control of Cabbage Seed Pod Weevil (*Ceutorhynchus assimilis*), treat when two or more Cabbage Seedpod Weevils may be found per plant on young pods.
- o For control of Aphids, treat about mid-growing season when insect populations reach economic thresholds.
- o For control of Flea Beetles, treat in early season when significant evidence of shot-holing by the adults becomes evident on rape seedlings.
- o For control of Diamondback Moth Larvae, treat about mid-growing season when populations reach economic thresholds and there is evidence of appreciable feeding beginning on rape flowers and/or pods.

A second application may be made if necessary. To protect pollinators apply only in the evening (this includes a period from 3 hours before sunset until midnight of the same day). Preharvest interval is 28 days; do not treat within 28 days of harvest). DO NOT graze treated fields or feed treated forage, ~~thrashing waste, or seed screenings~~ to livestock.

ACCEPTED
JUN - 8 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4787-15

Cheminova Agro A/S
P. O. Box 9
DK-7620
Lemvig, Denmark
Telephone: (+45) 97 83 4100

