

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

4787-23

11/01/2007

SUPPLEMENTAL LABELING

ACCEPTED

NOV - 1 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 4787-23



EPA Reg. No. 4787-23

Glyphos® X-TRA

Preharvest and Spot-Treatments of Weeds in Dry Peas, Lentils, and Chickpeas

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of application. All applicable directions and precautions in the Glyphos® X-TRA herbicide label must be followed.

See the "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the Glyphos® X-TRA herbicide label for essential product performance information.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

BROADCAST SPRAY: This product may be applied as an over the top broadcast spray to control labeled weeds prior to the harvest of dry peas, lentils, and chickpeas. Apply up to 24 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

PRECAUTION RESTRICTIONS:

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Glyphos® X-TRA herbicide label.
- Preharvest application is not recommended for peas, lentils, or chickpeas grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

SPOT TREATMENTS:

This product may be applied as spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry peas, lentils, and chickpeas. Apply up to 20 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a handheld sprayer. For best results, applications should be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

PRECAUTION RESTRICTIONS:

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Glyphos® X-TRA herbicide label.
- Preharvest application is not recommended for peas, lentils, or chickpeas grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

2/2

IMPORTANT

BEFORE USING GLYFOS X-TRA, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Read the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES on the Section 3 Federal product label.

This labeling must be in the possession of the user at the time of pesticide application.

Glyfos® is a registered trademark of Cheminova

Cheminova Inc.
1700 Route 23, Suite 300
Wayne, NJ 07470
www.cheminova.us.com

8/15/2007

NEXT

LABEL


4787-23

11/01/2007

1/2



SUPPLEMENTAL LABELING



Glyphos X-TRA

(EPA Reg. No. 4787-23)

Preharvest and Spot-Treatments of Weeds in Dry Beans

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Glyphos-XTRA herbicide for essential product performance information.

DIRECTIONS FOR USE

See the "APPLICATION EQUIPMENT AND TECHNIQUES" and "MIXING" section of the Glyphos X-TRA herbicide label booklet for more information.

To minimize drift, use nozzles that emit medium to large-sized droplets. The likelihood of drift occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.

BROADCAST SPRAY: This product may be applied as an over-the-top broadcast spray to control labeled weeds prior to the harvest of dry beans. Apply up to 22 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

PRECAUTION RESTRICTIONS:

- Apply at least 7 days before harvest.
- Only one application per year may be made; do not combine a pre-harvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Glyphos X-TRA herbicide label.
- Preharvest application is not recommended for dry beans grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat cowpeas, since these are considered to be grown as livestock feed.

SPOT TREATMENTS

2/2

Use Instructions: This product may be applied as spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry beans. Apply up to 22 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a handheld sprayer. For best results, applications should be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

PRECAUTION RESTRICTIONS:

- Apply at least 7 days before harvest.
- Only one application per year may be made; do not combine a pre-harvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Glyphos X-TRA herbicide label.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat cowpeas, since these are considered to be grown as livestock feed.

IMPORTANT

BEFORE USING GLYFOS X-TRA, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Read the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES on the Section 3 Federal product label.

This labeling must be in the possession of the user at the time of pesticide application.

Glyphos® is a registered trademark of Cheminova

Cheminova Inc.
1700 Route 23, Suite 300
Wayne, NJ 07470
www.cheminova.us.com

9/12/2007