

4787-34

04/28/2000

1/25

**Glyfos® AQ
Aquatic Herbicide**

For use in Aquatic and Other Non-Crop Sites.

**AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY
ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE
INJURY OR DESTRUCTION MAY RESULT.**

Carefully follow detailed instructions in label booklet.

Read the entire label before using this product

Use only according to label instructions

Read "DISCLAIMER" before buying or using. If terms are not acceptable, return
at once unopened.

**Keep Out Of Reach Of Children
CAUTION**

ACTIVE INGREDIENT:

* Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt	53.8 %
INERT INGREDIENTS:	46.2 %
TOTAL:	100.0 %

* Contains 651 grams per liter or 5.4 pounds per U. S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt.
Equivalent to 480 grams per liter or 4 pounds per U. S. gallon of the acid, glyphosate.

Manufactured by
CHEMINOVA AGRO A/S
P.O. Box 9, Lervig, Denmark


Authorized Representative:
Cheminova, Inc.
1700 Route 23, Oak Hill Park
Wayne, NJ 07470

® GLYFOS is a registered trademark of Cheminova Agro A/S

EPA Reg. No. 4787-XX

EPA Est. No.: _____

NET CONTENTSGallons (Liters)



DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Cheminova. All such risks shall be assumed by the user.

Cheminova warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the Directions for Use set forth in the Complete Directions for Use label booklet ("Directions"), subject to the risks referred to above.

Any damage arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

Cheminova makes no other express or implied warranty including any other express or implied warranty of FITNESS or MERCHANTABILITY.

The sale of this product does not include a license under any patent owned by Cheminova.

**PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals**

Keep Out Of Reach Of Children

CAUTION

- Harmful if absorbed through the skin. Avoid contact with eyes, skin or clothing.
- Avoid breathing spray mist or vapors.
- Remove contaminated clothing and wash before reuse.
- Wash thoroughly with soap and water after handling.

FIRST AID

- IF INHALED:** Remove individual to fresh air. Get medical attention if breathing difficulty develops.
- IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

IN CASE OF A MEDICAL EMERGENCY, CALL TOLL FREE DAY OR NIGHT, 1-800-228-5635, EXT. 133

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters. Treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss can cause fish suffocation.

In case of spill or leak, soak up with an absorbent and remove to a landfill.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED

STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or tribe, consult the agency responsible for pesticide regulations.

FOR MORE INFORMATION, CALL TOLL-FREE 1-800-548-6113

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: STORE ABOVE 10° F (-12° C) TO KEEP PRODUCT FROM CRYSTALIZING. Crystals will settle to the bottom. If allowed to crystalize, place in a warm room (≥ 68° F or 20° C) for several days to allow crystals to redissolve, then shake well before using.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State, or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container.

FOR BULK CONTAINERS: Triple rinse emptied bulk container. Then offer for recycling or reconditioning, or dispose of in a manner approved by state and local authorities.

FOR MINI-BULK REFILLABLE CONTAINERS: Do not reuse container, except for refill in accordance with a valid Cheminova Repackaging or Toll Repackaging Agreement. If not refilled or returned to an authorized repackaging facility, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR ALL OTHER NON-RETURNABLE/REFILLABLE CONTAINERS: Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This product, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of many herbaceous or woody plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "WEEDS CONTROLLED" section of this label. Unemerged plants arising from unattached

underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the recommended range when weed growth is heavy or dense.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

When this product comes into contact with soil, on the soil surface or as soil particles suspended in water, it is bound tightly to the soil particles. When used according to the label, once bound to soil particles, the product is no longer available for uptake by plants, and will not harm off-site vegetation where roots grow into the treatment area or if soil is transported off-site. When used according to the label, the strong affinity of this product to soil prevents leaching out of the soil and entering ground water. The affinity remains until the product is degraded both aerobically, and anaerobically, primarily by soil microorganisms.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

The maximum use rate stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate(s).

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift.

AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animal or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. DO NOT APPLY WHEN WIND OR OTHER CONDITIONS FAVOR DRIFT. HAND-GUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS. NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

MIXING

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of this product (see the "DIRECTIONS FOR USE" and "WEEDS CONTROLLED" sections of this label) near the end of the filling process and mix well. Remove hose from tank immediately after filling to avoid siphoning back into the carrier source. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in in-line strainer or nozzles should be 50 mesh or larger. Use correct nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

When using this product, add 2 or more quarts of non-ionic surfactant per 100 gallons of spray solution. Use a non-ionic surfactant labeled for use with herbicides. The surfactant must contain 50% or more of active ingredient. Always read and follow the surfactant manufacturer's label instructions. Surfactants should not be used in excess of 1 quart/A in broadcast application.

Colorants or marking dyes approved for use with herbicides may be added to spray mixtures of this product, but may reduce performance, especially at low rates. Always read and follow the colorant manufacturer's label directions.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water. Dispose of rinsate according to labeled use or disposal instructions.

Carefully observe all cautionary statements and other information on the surfactant label.

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

Aerial - Fixed Wing and Helicopter

Broadcast Spray

Controlled Droplet Applicator (CDA) - Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

Hand-Held and High-Volume Spray Equipment - Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage. Use only coarse sprays.

See the appropriate part of this section for specific instructions and rates of application.

AERIAL EQUIPMENT

Use the recommended rates of this herbicide in 3 to 20 gallons of water per acre unless otherwise specified on this label. See the "WEEDS CONTROLLED" section of this label for specific rates. Aerial applications of this product may be made as specifically stated on this label. Refer to the individual use area sections of this label for recommended volumes and application rates.

AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements

and all other information appearing on the additive label.

Ensure uniform application — To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

SPRAY DRIFT MANAGEMENT

DRIFT MAY CAUSE DAMAGE TO ANY OTHER VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

- 1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- 2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

Importance of Droplet Size - The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion sections of this label).

Controlling Droplet Size - Volume: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure: Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy protection. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles: Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation: Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type: Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment - When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind - Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity - When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversion - Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves literally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that move upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas - The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

BROADCAST EQUIPMENT

For control of annual or perennial weeds listed on this label using broadcast equipment - Use the recommended rates of this product in 3 to 30 gallons of water per acre as a broadcast spray unless otherwise specified on this label. See the "WEEDS CONTROLLED" section of this label for specific rates. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. For better results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

CONTROLLED DROPLET APPLICATION (CDA)

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre. Do not apply as a fine mist. Use coarse sprays only.

For the control of labeled annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 MPH (1 quart per acre). For the control of labeled perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and walking speed of 0.75 MPH (2 to 4 quarts per acre).

Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

HAND-HELD and HIGH-VOLUME EQUIPMENT

Use coarse sprays only.

Mix this product in clean water and apply to foliage of vegetation to be controlled. Applications should be made on a spray-to-

wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. For use in knapsack sprayers, it is suggested that the recommended amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

For control of annual weeds listed on this label, apply a 3/4 to 2 percent solution of this product plus nonionic surfactant and apply to foliage of vegetation to be controlled.

When using application methods which result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 8 percent solution for woody brush and trees. If a straight-stream nozzle is used, start the application at the top of the target vegetation and spray from top to bottom in a "zig-zag" motion. Ensure that at least 50% of the leaf surface is contacted by the spray.

Small open-branched trees need only be treated from one side. If foliage is thick or there are multiple root sprouts, application must be made from several sides to ensure adequate coverage.

Recommended mixtures are shown in the following table:

Amount of GLYFOS AQ Herbicide						
concentration	3/4%	1%	1 1/4%	1 1/2%	5%	8%
Volume						
1 gallon	1 oz	1 1/3 oz	1 2/3 oz	2 oz	6 2/3 oz	10 1/4 oz
25 gallons	1 1/2 pint	1 qt	1 1/4 qt	1 1/2 qt	5 qt	2 gal
100 gallons	3 qt	1 gal	1 1/4 gal	1 1/2 gal	5 gal	8 gal
2 tablespoons = 1 fluid ounce						

WEEDS CONTROLLED

ANNUAL WEEDS

Apply to actively growing grass and broadleaf weeds. Allow at least 3 days after treatment before disturbing vegetation. After this period, weeds may be mowed tilled or burned. To prevent seed production, applications should be made prior to seedhead formation.

Broadcast Application: Use 1 1/2 pints of this product per acre plus 2 or more quarts of a non-ionic surfactant per 100 gallons of spray solution when weeds are less than 6 inches tall. If weeds are greater than 6 inches tall, use 2 1/2 pints of this product per acre plus 2 or more quarts of nonionic surfactant per 100 gallons of spray solution.

Hand-held High-volume Application: Use a 3/4 % solution of this product in water plus 2 or more quarts of nonionic surfactant per 100 gallons of spray solution and apply to foliage of vegetation to be controlled.

This product will control the following annual weeds when applied as directed. Annual weeds will continue to germinate from seed throughout the growing season. Repeat treatments will be necessary to control later germinating seeds.

**balsamapple <i>Momordica charantia</i>	fiddleneck <i>Amsinckia</i> spp.
barley <i>Hordeum vulgare</i>	flaxleaf fleabane <i>Conyza bonariensis</i>
barnyardgrass <i>Echinochloa crus-galli</i>	fleabane <i>Erigeron</i> spp.
Bassia, fivehook <i>Bassia hyssopifolia</i>	foxtail <i>Setaria</i> spp.
bluegrass, annual <i>Poa annua</i>	foxtail, Carolina <i>Alopecurus carolinianus</i>
bluegrass, bulbous <i>Poa bulbosa</i>	groundsel, common <i>Senecio vulgaris</i>
bromegrass <i>Bromus</i> spp.	horseweed/ marestail <i>Conyza canadensis</i>
buttercup <i>Ranunculus</i> spp.	kochia <i>Kochia scoparia</i>
cheat <i>Bromus secalinus</i>	lambsquarters, common <i>Chenopodium album</i>
chickweed, mouseear <i>Cerastium vulgatum</i>	lettuce, prickly <i>Lactuca serriola</i>
cocklebur <i>Xanthium strumarium</i>	morningglory <i>Ipomoea</i> spp.
corn <i>Zea mays</i>	mustard, blue <i>Chorispora tenella</i>
crabgrass <i>Digitaria</i> spp.	mustard, tansy <i>Descurainia pinnata</i>
dwarf dandelion <i>Krigia cespitosa</i>	mustard, tumble <i>Sisymbrium altissimum</i>
false flax, smallseed <i>Camelina microcarpa</i>	mustard, wild <i>Sinapis arvensis</i>

oats, wild *Avena fatua*

panicum *Panicum spp.*

pennycress, field *Thlaspi arvensis*

pigweed, redroot *Amaranthus retroflexus*

pigweed, smooth *Amaranthus hybridus*

ragweed, common *Ambrosia artemisiifolia*

ragweed, giant *Ambrosia trifida*

rocket, London *Sisymbrium irio*

rye *Secale cereale*

*ryegrass Italian *Lolium multiflorum*

sandbur, field *Cenchrus spp.*

shattercane *Sorghum bicolor*

shepherdspurse *Capsella bursa-pastoris*

signalgrass, broadleaf *Brachiaria platyphylla*

smartweed, Pennsylvania *Polygonum pennsylvanicum*

sowthistle, annual *Sonchus oleraceus*

*Spanishneedles *Bidens bipinnata*

stinkgrass *Eragrostis cilianensis canadensis*

sunflower *Helianthus annuus*

thistle, Russian *Salsola kali*

spurry, umbrella *Holosteum umbellatum*

velvetleaf *Abutilon theophrasti*

wheat *Triticum aestivum*

witchgrass *Panicum capillare*

* Apply 3 pints of this product per acre

** Apply with hand-held equipment only

† Except glyphosate tolerant varieties and hybrids

11/25

PERENNIAL WEEDS

Apply this product as follows to control or destroy most vigorously growing perennial weeds. Unless otherwise directed, allow at least 7 days after application before disturbing vegetation.

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages. Fall treatments must be applied before a killing frost.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed.

Add 2 or more quarts of nonionic surfactant per 100 gallons of spray solution to the rates of this product given in this list. Refer to the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label for specific uses and applications instructions.

When applied as recommended under the conditions described, this product WILL CONTROL the following PERENNIAL WEEDS:

- | | |
|---|---|
| Alfalfa <i>Medicago sativa</i> | Dandelion <i>Taraxacum officinale</i> |
| Alligatorweed* <i>Alternanthera philoxeroides</i> | Dock, curly <i>Rumex crispus</i> |
| Anise/ Fennel <i>Foeniculum vulgare</i> | Dogbane, hemp <i>Apocynum cannabinum</i> |
| Artichoke, Jerusalem <i>Helianthus tuberosus</i> | Fescues <i>Festuca spp.</i> |
| Bahiagrass <i>Paspalum notatum</i> | Fescue, tall <i>Festuca arundinacea</i> |
| Bermudagrass <i>Cynodon dactylon</i> | Guineagrass <i>Panicum maximum</i> |
| Bermudagrass, water (knotgrass) <i>Paspalum distichum</i> | Hemlock, poison <i>Conium maculatum</i> |
| Bindweed, field <i>Convolvulus arvensis</i> | Horsenettle <i>Solanum carolinense</i> |
| Bluegrass, Kentucky <i>Poa pratensis</i> | Horseradish <i>Armoracia rusticana</i> |
| Blueweed, Texas <i>Helianthus ciliaris</i> | Ice plant <i>Mesembryanthemum crystallinum</i> |
| Brackenfern <i>Pteridium aquilinum</i> | Johnsongrass <i>Sorghum halepense</i> |
| Bromegrass, smooth <i>Bromus inermis</i> | Kikuyugrass <i>Pennisetum clandestinum</i> |
| Canarygrass, reed <i>Phalaris arundinacea</i> | Knapweed <i>Centaurea repens</i> |
| Cattail <i>Typha spp</i> | Lantana <i>Lantana camara</i> |
| Clover, red <i>Trifolium pratense</i> | Lespidiza, common, serices <i>Lespidiza striata</i> ,
<i>Lespidiza cuneata</i> |
| Clover, white <i>Trifolium repens</i> | Loosestrife, purple <i>Lythrum salicaria</i> |
| Cogongrass <i>Imperata cylindrica</i> | Lotus, American <i>Nelumbo lutea</i> |
| Cordgrass <i>Spartina spp.</i> | Maidencane <i>Panicum hamatomon</i> |
| Cutgrass, giant* <i>Zizaniopsis miliacea</i> | Milkweed <i>Asclepias spp.</i> |
| Dallisgrass <i>Paspalum dilatatum</i> | Muhly, wirestem <i>Muhlenbergia frondosa</i> |

Mullein, common *Verbascum thapsus*

Napiergrass *Pennisetum purpureum*

Nightshade, silverleaf *Solanum elaeagnifolium*

Nutsedge, purple, yellow *Cyperus rotundus*
Cyperus esculentus

Orchardgrass *Dactylis glomerata*

Pampasgrass *Cortaderia jubata*

Paragrass *Brachiaria mutica*

Phragmites** *Phragmites spp.*

Quackgrass *Elymus repens*

Reed, giant *Arundo donax*

Ryegrass, perennial *Lolium perenne*

Smartweed, swamp *Polygonum coccineum*

Spatterdock *Nuphar luteum*

Starthistle, yellow *Centaurea solstitialis*

Sweet potato, wild* *Ipomoea pandurata*

Thistle, artichoke *Cynara cardunculus*

Thistle, Canada *Cirsium arvense*

Timothy *Phleum pratense*

Torpedograss* *Panicum repens*

Tules, common *Scirpus acutus*

Vaseygrass *Paspalum urvillei*

Velvetgrass *Holcus spp.*

Waterhyacinth *Eichornia crassipes*

Waterlettuce *Pistia stratiotes*

Waterprimrose *Ludwigia spp.*

Wheatgrass, western *Agropyron smithii*

* Partial Control

** Partial control, Southeastern States, see specifics below.

See "DIRECTIONS FOR USE" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for labeled uses and specific application instructions.

Alligatorweed - Apply 6 pints of this product per acre or apply a 1 1/4 percent solution with hand-held equipment to provide partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain such control.

Bermudagrass - Apply 7 1/2 pints of this product per acre as a broadcast spray, or as a 1 1/2 percent solution with hand-held equipment. Treat when bermudagrass is actively growing and seed heads are present. Retreatment may be necessary to maintain control.

Bindweed, field; Silverleaf nightshade; Texas Blueweed: Apply 6 to 7 1/2 pints of this product per acre as a broadcast spray west of the Mississippi River and 4 1/2 to 6 pints per acre east of the Mississippi River. With hand-held equipment, use a 1 1/2 percent solution. Apply when target plants are actively growing, and are at or beyond full bloom. For silverleaf nightshade, best results can be obtained when application is made after berries are formed. Do not treat when weeds are under drought stress. New leaf development indicates active growth. For best results apply in late summer or fall.

Brackenfern - Apply 4 1/2 to 6 pints of this product per acre as a broadcast spray or as a 3/4 to 1 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.

Cattail: Apply 4 1/2 to 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when the plants are actively growing and are at or beyond the early to full bloom stage of growth. Best results are achieved when application is made during the summer or fall months.

Cogongrass - Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray. Apply when Cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Cordgrass: Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment. Schedule applications in order to allow 6 hours before treated plants are covered by tidewater. The presence of silt or debris on the cordgrass will reduce performance. It may be necessary to wash target plants before application to improve uptake.

Cutgrass, giant: Apply 6 pints of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment to provide partial control of giant cutgrass. Repeat applications will be required to maintain control, especially where vegetation is partially submerged. Allow regrowth to the 7 to 10 leaf stage before retreatment.

Dogbane, hemp; knapweed; horseradish: Apply 6 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing, and most have reached the late bud to flower stage of growth. For best results apply in late summer or fall.

Fescue, tall - Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Apply when the plants are actively growing, and most have reached the boot to head stage of growth. When applied prior to boot stage, control may be reduced.

Guineagrass: Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when plants are actively growing and most have reached at least the 7-leaf stage of growth.

Johnsongrass; Kentucky bluegrass; Smooth brome grass; Reed canarygrass; Orchardgrass; Perennial ryegrass; Timothy; Western wheatgrass: Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing, and most have reached the boot to head stage of growth. When applied prior to boot stage, control may be reduced. In fall, apply before plants have turned brown.

Lantana: Apply this product as a 3/4 to 1 percent solution with hand-held equipment. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher rate for plants that have reached the woody stage of growth.

Loosestrife, purple: Apply 4 pints of this product per acre as a broadcast spray or as a 1 to 1 1/2 percent solution with hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost.

Lotus, American: Apply 4 pints of this product as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost. Repeat treatments may be required to control regrowth from seeds or underground parts of plants.

Maidencane; Paragrass: Apply 6 pints of this product as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Repeat treatments will be required, especially with plants that are partially submerged in water. Under these conditions, allow regrowth to the 7 to 10 leaf stage of growth before retreatment.

Milkweed, common: Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when plants are actively growing and have reached the late bud to flower stage of growth.

Nutsedge, purple/yellow: Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment to control existing plants or immature nutlets attached to treated plants. Apply when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control.

Pampasgrass: Apply a 1 1/2 percent solution of this product with hand-held equipment when plants are actively growing.

Phragmites: For partial control of phragmites in Florida and the counties of other states bordering on the Gulf of Mexico, apply 7 1/2 pints per acre of this product as a broadcast spray, or as a 1 1/2 percent solution with hand-held equipment. In other areas of the US, apply 4 to 6 pints per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment for partial control. For best results treat during late summer or fall months when plants are actively growing and in full bloom. Due to the dense nature of the vegetation which may prevent good spray coverage and uneven stages of growth, repeat treatments may be necessary to maintain control. Visual symptoms will be slow to develop.

Quackgrass; Kikuyugrass; Wirestem muhly: Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment when most quackgrass or wirestem muhly is at least 8 inches high (3 to 4 leaf stage of growth) and actively growing. Allow 3 or 4 more days after application before tillage.

Giant reed; Iceplant: For control of giant reed and iceplant, apply a 1 1/2 percent solution of this product with hand-held equipment when plants are actively growing. For giant reeds best results are obtained when application is made in late summer to fall.

Spatterdock: Apply 6 pints per acre of this product as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when most of the plants are in full bloom. For best results apply in late summer or fall.

Wild Sweet Potato: Apply as a 1 1/2 percent solution with hand-held equipment. Apply to actively growing weeds at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment.

Thistle, Canadal Artichoke: Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held for Canada thistle. To control artichoke thistle, apply a 2 percent solution as a spray-to-wet application. Apply when plants are actively growing and are at or beyond the bud stage of growth.

Torpedograss: Apply 6 to 7 1/2 pints per acre of this product as a broadcast spray or as 3/4 to 1 1/2 percent solution with hand-held equipment to provide partial control of torpedograss. Use the lower rates under terrestrial conditions, and the higher rates under partially submerged or floating-mat conditions. Repeat treatment will be required to maintain control.

Tules, common: Apply this product as a 1 1/2 percent solution with hand-held equipment. Apply to actively growing plants at or beyond the seed head stage of growth. After application visual symptoms will be slow to appear and may not appear for 3 or more weeks.

Waterhyacinth: Apply 5 to 6 pints of this product per acre as a broadcast spray or as a 3/4 to 1 percent solution with hand-held equipment. Apply when plants are actively growing and at or beyond stage of growth. After application, visual symptoms may require 3 or more weeks to appear with complete necrosis and decomposition usually occurring within 60 to 90 days. Use the higher rates when more rapid visual effects are required.

Waterlettuce: For control, apply a 3/4 to 1 percent solution of this product with hand-held equipment to actively growing plants. Use the higher rates where infestations are heavy. Best results are obtained from mid-summer through winter applications. Spring application may require retreatment.

Waterprimrose: Apply this product as a 3/4 percent solution using hand-held equipment to plants that are actively growing at or beyond the bloom stage of growth, but before fall color changes occur. Thorough coverage is necessary for best control.

Other perennials listed on this label: Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing, and most have reached early head or early bud stage of growth.

WOODY BRUSH AND TREES

When applied as recommended under the conditions described, this product CONTROLS or PARTIALLY CONTROLS the following woody brush, plants and trees:

Alder
Alnus spp.

Ash*
Fraxinus spp.

Aspen, quaking
Populus tremuloides

Bearmat (Bearclover)
Chamaebatia foliolosa

Birch
Betula spp.

Blackberry
Rubus spp.

Broom:
French
Cytisus monspessulanus
Scotch
Cytisus scoparius

Buckwheat, California*
Eriogonum fasciculatum

Cascara*
Rhamnus purshiana

Catsclaw*
Acacia greggi

Ceanothus
Ceanothus spp.

Chamise
Adenostoma fasciculatum

Cherry:
Bitter
Prunus emarginata
Black
Prunus serotina
Pin
Prunus pensylvanica

Coyote brush
Baccharis consanguinea

Creeper, Virginia*
Parthenocissus quinquefolia

Dewberry
Rubus trivialis

Dogwood
Cornus spp.

Elderberry
Sambucus spp.

Elm*
Ulmus spp.

Eucalyptus, bluegum
Eucalyptus spp.

Hasardia*
Haplopappus squamosus

Hawthorn
Crataegus spp.

Hazel
Corylus spp.

Hickory
Carya spp.

Holly, Florida/
Brazilian Peppertree
Schinus terebinthifolius

Honeysuckle
Lonicera spp.

Hornbeam, American
Carpinus caroliniana

Kudzu
Pueraria lobata

Locust, black*
Robinia pseudoacacia

Manzanita
Arctostaphylos spp.

Maple:
Red**
Acer rubrum
Sugar
Acer saccharum
Vine*
Acer circinatum

Monkey Flower*
Mimulus guttatus

Oak
Black*
Quercus velutina —
Northern Pine
Quercus palustris
Post
Quercus stellata
Red
Quercus rubra
Southern Red
Quercus falcata
White*
Quercus alba

Persimmon*
Diospyros spp.

Poison Ivy
Rhus radicans —

Poison Oak
Rhus toxicodendron

Poplar, yellow* (Tulip tree)
Liriodendron tulipifera

Prunus
Prunus spp.

Raspberry
Rubus spp.

Redbud, eastern
Cercis canadensis

Rose, multiflora
Rosa multiflora

Russian olive
Elaeagnus angustifolia

Sage: black, white
Salvia spp.

Sagebrush, California
Artemisia californica
Salmonberry
Rubus spectabilis

Salt cedar
Tamarix spp.

Saltbush, Sea myrtle
Baccharis halimifolia

Tallowtree, Chinese
Sapium sebiferum

Sassafras
Sassafras spp.

Sourwood*
Oxydendrum arboreum Sumac:

Thimbleberry
Rubus parviflorus

Poison*
Rhus vernix

Tobacco, tree*
Nicotiana glauca

Smooth*
Rhus glabra

Winged*
Rhus copallina

Trumpet creeper
Campsis radicans

Sweetgum
Liquidambar styraciflua

Waxmyrtle, southern*
Myrica cerifera

Swordfern*
Polystichum munitum

Willow
Salix spp.

* Partial control

** See below for control or partial control instructions.

NOTE: If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stages of growth.

Apply this product when plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering.

) Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See "DIRECTIONS FOR USE", and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for labeled uses and specific application instructions.

Applied as a 5 to 8 percent solution as a directed application as described in the "HAND HELD AND HIGH VOLUME EQUIPMENT" section, this product will control or partially control all the species listed in this section of this label. Use the higher rate for dense stands and larger woody brush and trees.

Apply this product as follows to control or partially control the following woody brush and trees:

Alder / Black berry / Dewberry / Honeysuckle / Post Oak / Raspberry - For control, apply 4 1/2 to 6 pints per acre of this product as a broadcast spray or as a 3/4 to 1 1/4 percent solution with hand-held equipment.

Aspen, quaking / Hawthorn / Trumpet creeper - For control, apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment.

Birch / Elderberry / Hazel / Salmonberry / Thimbleberry - For control, apply 3 pints per acre of this product as a broadcast spray or as a 3/4 percent solution with hand-held equipment.

Broom: French, Scotch - For control, apply a 1 1/4 to 1 1/2 percent solution with hand-held equipment.

Buckwheat, California / Hasardia / Monkey Flower / Tobacco, tree - For partial control of these species, apply a 3/4 to 1 1/2 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Catsclaw - For partial control, apply as a 1 1/4 to 1 1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Cherry: Bitter, Black, Pin / Oak, Southern Red / Sweet Gum / Prunus - For control, apply 3 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 to 1 1/2 percent solution with hand-held equipment.

Coyote Brush - For control, apply a 1 1/4 to 1 1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Dogwood / Hickory / Salt cedar - For partial control, apply a 1 to 2 percent solution of this product with hand-held equipment or 6 to 7 1/2 pints per acre as a broadcast spray.

Eucalyptus, bluegum - For control of eucalyptus resprouts, apply a 1 1/2 percent solution of this product with hand-held equipment when resprouts are 6 to 12 feet tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application to drought-stressed plants.

Holly, Florida / Waxmyrtle, southern - For partial control, apply this product as a 1 1/2 percent solution with hand-held equipment.

Kudzu - For control, apply 6 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.

Maple, red - For control, apply as a 3/4 to 1 1/4 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 7 1/2 pints of this product per acre as a broadcast spray.

Maple, sugar / Oak, northern pin / Oak, red - For control, apply as a 3/4 to 1 1/4 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Poison Ivy / Poison Oak - For control, apply 6 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

Rose, multiflora - For control, apply 3 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects.

Sage, black / Sagebrush, California / Chamise / Tallowtree, Chinese - For control of these species, apply a 3/4 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Saltbush, Sea myrtle - For control, apply this product as a 1 percent solution with hand-held equipment.

Tan oak resprouts - For suppression or partial control, apply a 2 percent solution of this product to resprouts less than 3 to 6 feet tall. Best results are obtained with fall applications.

Willow - For control, apply 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment.

Other Woody Brush and Trees listed on this label - For partial control, apply 3 to 7 1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1 1/2 percent solution with hand-held equipment.

AQUATIC AND OTHER NONCROP SITES

When applied as directed and under the conditions described in the "Weeds Controlled" section in this label this product will control or partially control the labeled weeds growing in the following industrial, recreational, and public areas or other similar aquatic and terrestrial sites.

Aquatic Sites - This product may be applied to emerged weeds in all bodies of fresh and brackish water which may be flowing, non-flowing, or transient. This includes lakes, rivers, streams, ponds, estuaries, rice levees, seeps, irrigation and drainage ditches, canals, reservoirs, wastewater treatment facilities, wildlife habitat restoration and management areas, and similar sites.

If aquatic sites are present in the non-crop area and are part of the intended treatment, read and observe the following directions:

This product does not control plants which are completely submerged or have a majority of foliage under water.

There is no restriction on the use of treated water for irrigation, recreation, or domestic purposes.

Consult local state fish and game agency and water control authorities before applying this product to public waters. Permits may be required to treat such water.

NOTE: Do not apply this product within 1/2 mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as lake, pond, or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis. These aquatic applications may be made **ONLY** in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.

For treatments after drawdown of water or in dry ditches, allow 7 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after drawdown to ensure application to actively growing weeds.

Floating mats of vegetation may require retreatment. Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash or by rainfall within 6 hours of application. Do not re-treat within 24 hours following the initial treatment.

Applications made to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in water. When making any bankside applications, do not overlap more than 1 foot into open water. Do not spray in bodies of water where weeds do not exist. The maximum application rate of 7 1/2 pints per acre must not be exceeded in any single broadcast application that is being made over water.

When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in fish kill.

21/25

Other Noncrop-Type Sites - This product may be used to control the listed weeds in terrestrial noncrop sites and/or in aquatic sites within these areas:

Airports
Golf Courses
Habitat Restoration & Management Areas
Highways & Roadsides
Industrial Plant Sites
Lumberyards
Parking Areas
Parks

Petroleum Tank Farms
Pipeline, Power, Telephone & Utility Rights-of-Way
Pumping Installations
Railroads
Schools
Similar Sites
Storage Areas

WILDLIFE HABITAT RESTORATION AND MANAGEMENT AREAS

This product is recommended for the restoration and/or maintenance of native habitat and in wildlife management areas.

Habitat Restoration and Maintenance - When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications may be made to allow recovery of native plant species, to open up water to attract waterfowl, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments may be made to selectively remove unwanted plants for habitat enhancement. For spot treatments, care should be exercised to keep spray off of desirable plants.

Wildlife Food Plots - This product may be used as a site preparation treatment prior to planting wildlife food plots. Apply as directed to control vegetation in the plot area. Any wildlife food species may be planted after applying this product, or native species may be allowed to reinfest the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling to allow for maximum effectiveness.

WIPER APPLICATIONS

For wick or wiper applications, mix 1 gallon of this product with 2 gallons of clean water to make a 33 percent solution. Addition of a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution is recommended.

Wiper applications can be used to control or suppress annual and perennial weeds listed on this label. In heavy weed stands, a double application in opposite directions may improve results. See the "Weeds Controlled" section in this label for recommended timing, growth stage and other instructions for achieving optimum results.

CUT STUMP APPLICATION

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of this product to freshly cut surface immediately after cutting. Delay in applying this product may result in reduced performance. For best results, trees should be cut during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will CONTROL, PARTIALLY CONTROL or SUPPRESS most woody brush and tree species, some of which are listed below:

Alder <i>Alnus spp.</i>	Poplar* <i>Populus spp.</i>
Coyote brush* <i>Baccharis consanguinea</i>	Reed, giant <i>Arundo donax</i>
Dogwood* <i>Cornus spp.</i>	Salt cedar <i>Tamarix spp.</i>
Eucalyptus <i>Eucalyptus spp.</i>	Sweet gum* <i>Liquidambar styraciflua</i>
Hickory* <i>Carya spp.</i>	Sycamore* <i>Platanus occidentalis</i>
Madrone <i>Arbutus menziesii</i>	Tan Oak <i>Lithocarpus densiflorus</i>
Maple* <i>Acer spp.</i>	Willow <i>Salix spp.</i>
Oak <i>Quercus spp.</i>	

* This product is not approved for this use on these species in the state of California.

INJECT AND FRILL APPLICATIONS

Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment which must penetrate into living tissue. Apply the equivalent of 1 ml of this product per 2 to 3 inches of trunk diameter. This is best achieved by applying 25 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying dilute material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as these, make frill or cut at an oblique angles so as to produce a cupping effect and use undiluted material. For best results, applications should be made during periods of active growth and full leaf expansion.

This treatment WILL CONTROL the following woody species:

Oak <i>Quercus spp.</i>	Sweet gum <i>Liquidambar styraciflua</i>
Poplar <i>Populus spp.</i>	Sycamore <i>Platanus occidentalis</i>

This treatment WILL SUPPRESS the following woody species:

Black gum* <i>Nyssa sylvatica</i>	Hickory <i>Carya spp.</i>
Dogwood <i>Cornus spp.</i>	Maple, red <i>Acer rubrum</i>

*This product is not approved for this use on this species in the state of California.

RELEASE OF BERMUDAGRASS OR BAHIAGRASS ON NONCROP SITES

RELEASE OF DORMANT BERMUDAGRASS AND BAHIAGRASS

When applied as directed, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Make applications to dormant bermudagrass or bahiagrass.

For best results on winter annuals, treat when weeds are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage.

WEEDS CONTROLLED

Rate recommendations for control or suppression of winter annuals and tall fescue are listed below.

Apply the recommended rates of this product in 10 to 25 gallons of water per acre plus 2 quarts nonionic surfactant per 100 gallons of total spray volume.

For the best recommendation for the mixture of weeds within your geographic area, contact your sales representative.

WEEDS CONTROLLED OR SUPPRESSED*

Note: C = Control
S = Suppression

WEED SPECIES	Glyfos® AQ FLUID OZ/ACRE					
	6	9	12	18	24	48
Barley, little <i>Hordeum pusillum</i>	S	C	C	C	C	C
Bedstraw, catchweed <i>Galium aparine</i>	S	C	C	C	C	C
Bluegrass, annual <i>Poa annual</i>	S	C	C	C	C	C
Chervil <i>Chaerophyllum tainturieri</i>	S	C	C	C	C	C
Chickweed, common <i>Stellaria media</i>	S	C	C	C	C	C
Clover, crimson <i>Trifolium incarnatum</i>	.	S	S	C	C	C
Clover, large hop <i>Trifolium campestre</i>	.	S	S	C	C	C
Fescue, tall <i>Festuca arundinaceae</i>	S	S
Geranium, Carolina <i>Geranium carolinanum</i>	.	.	S	S	C	C
Henbit <i>Lamium amplexicaule</i>	.	S	C	C	C	C
Ryegrass, Italian <i>Lolium multiflorum</i>	.	.	S	C	C	C
Speedwell, corn <i>Veronica arensis</i>	S	C	C	C	C	C
Vetch, common <i>Vicia sativa</i>	.	.	S	C	C	C

* These rates apply only to sites where an established competitive turf is present.

RELEASE OF ACTIVELY GROWING BERMUDAGRASS

NOTE: USE ONLY ON SITES WHERE BAHIAGRASS OR BERMUDAGRASS ARE DESIRED FOR GROUND COVER AND SOME TEMPORARY INJURY OR YELLOWING OF THE GRASSES CAN BE TOLERATED.

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "Weeds Controlled" section on this label, and suppression or partial control of certain perennial weeds.

For control or suppression of those annual species listed in this label, use 3/4 to 2 1/4 pints of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre, plus 2 quarts of nonionic surfactant per 100 gallons of total spray volume. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use higher rate as size of plants increases or as they approach or seedhead formation.

Use the higher rate for partial control or longer-term suppression of the following perennial species. Use lower rates for shorter-term suppression of growth.

Bahiagrass
Dallisgrass
Fescue (tall)

Johnsongrass**
Trumpetcreeper*
Vaseygrass

- * Suppression at the higher rate only.
- ** Johnsongrass is controlled at the higher rate.

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may result.

BAHIAGRASS SEEDHEAD AND VEGETATIVE SUPPRESSION

When applied as directed in the "Noncrop Sites" section in this label, this product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications.

Apply this product 1 to 2 weeks after full greenup of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 5 fluid ounces per acre of this product plus 2 quarts of an approved nonionic surfactant per 100 gallons of total spray volume in 10 to 25 gallons of water per acre.

Sequential applications of this product plus nonionic surfactant by total spray volume may be made at approximately 45 day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence.

Apply no more than 2 sequential applications per year. As a first sequential application, apply 3 fluid ounces of this product per acre plus nonionic surfactant. A second sequential application of 2 to 3 fluid ounces per acre plus nonionic surfactant may be made approximately 45 days after the last application.

ANNUAL GRASS GROWTH SUPPRESSION

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 3 to 4 ounces of this product in 10 to 40 gallons of spray solution per acre. Mix 2 quarts of a nonionic surfactant per 100 gallons of spray solution. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

© Cheminova, 1999