

4787-46

04/13/2004

S748095- 1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

APR 13 2004

Ms. Diane Allemang
Director, Regulatory Affairs
Cheminova Agro A/S
c/o Cheminova Inc.
600 13th Street, N.W.
Washington, DC 20005

Dear Ms Allemang:

Subject: Updated Label
Atrapa 8E
EPA Reg. No. 4787-46 (Formally, 1812-408)
Your Application of October 10, 2003

The labeling amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, As Amended (FIFRA), is acceptable, provided that you submit two copies of your final printed label incorporating the correction cited below before you release the product for shipment.


Reinstate the statement below as it appears on the registered label dated February 10, 1999.

Ultra-low volume (ULV) application is limited to the crops alfalfa, clover, pasture and range grass, grass and grass hay, grain crops (barley, corn, oats, rye & wheat), beans, rice and nonagricultural lands.

If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended label constitutes acceptance of this condition.

A stamped copy of the label is enclosed for your records.

Sincerely,


Marilyn A. Mautz
Biologist
Insecticide-Rodenticide Branch
Registration Division (7504C)

10/10/03

Atrapa® 8E

APR 13 2004

g408s03a

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

Active Ingredient

Malathion: O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate ^{4787-46 (Formerly 1812-408)} 81.43%

Inert Ingredients 18.57%

Total 100.00%

This Product Contains 8 Pounds of Malathion per Gallon

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID Organophosphate	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies involving this product, call toll free 1-888-324-7598.</p>	
<p>NOTE TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine.</p>	
<p>See Label for Additional Precautions and Directions for Use.</p>	

Manufactured for:
Griffin L.L.C.
Valdosta, GA 31601

EPA Reg. No. 1812-408
EPA Est. No. _____

Net Contents _____

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through the skin. Causes eye irritation. Avoid breathing spray mist. Do not get in eyes, on skin or clothing.

~~**STATEMENT OF PRACTICAL TREATMENT**~~

~~Organophosphate~~

~~If swallowed, do not induce vomiting. Get medical attention. In case of contact, immediately flush eyes and skin with plenty of water for at least 15 minutes. Call a physician.~~

~~**NOTE TO PHYSICIAN:** This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine.~~

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates and aquatic life stages of amphibians. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read directions carefully. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and the handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. In case of spill or leak, soak up with sand, earth or synthetic absorbent and dispose as waste.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill ~~or by other procedures approved by state and local authorities~~ or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Pour specified amount of this product into nearly filled spray tank. Add balance of water to filled tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of this mixture has proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION: (The actual rate required to provide thorough, uniform coverage varies with plant growth at time of application. Except as specified for certain uses, the following rates are therefore intended to cover a broad range of conditions.)

Crop	Gallons of Water per Acre	
	Aircraft	Ground
Vegetable and Field Crops	1-20	5-125
Orchard, Grapes	5-25	
Orchard Crops (see exceptions below)		300-800
Citrus		500-3000
Grapes		100-200

6/12

RECOMMENDATIONS

Unless otherwise indicated, dosages are given in pints of Atrapa 8E per acre in sufficient water to provide thorough coverage of vegetable and field crops, and in pints of Atrapa 8E per 100 gallons of water for thorough coverage sprays of fruit crops. Begin applications when insects first appear and repeat at 7 to 10 day intervals as needed to maintain control, but observe use limitations given for specific crops. Maximum permissible rate per acre, expressed as Atrapa 8E, given in parenthesis () after each crop claim.

~~Ultra-low volume (ULV) application is limited to the crops alfalfa, clover, pasture and range grass, grass and grass hay, grain crops (barley, corn, oats, rye, & wheat), beans, rice, and nonagricultural lands.~~

CROP USES

TREE FRUIT CROPS

APRICOT: Aphids, Codling Moth, European Lecanium Scale, Orange Tortrix, Soft Brown Scale - 1¼ pints. Do not apply within 7 days of harvest. (10 pints)

CHERRY: Black Cherry Aphid, Bud Moth, Cherry Fruit Fly, Fruit Tree Leaf Roller, Lesser Peach Tree Borer (apply to trunks and scaffold limbs at 21-day intervals beginning with emergence), **Forbes and San Jose Scales** (apply 3 weeks before and immediately after harvest) - 1 pint. Do not apply within 3 days of harvest. **Caution NOTE:** May cause injury on certain varieties of sweet cherry in the Northeast. (8 pints)

CITRUS (Orange, Grapefruit, Kumquat, Lemon, Lime, and Tangerine): California Red Scale, Yellow Scale, Purple Scale, Black Scale, (Single and Off-Brooded) Soft Scale, Thrips, Citricola Scale, Aphids - 1 pint. Do not apply within 7 days of harvest. Make no application when trees are in bloom. **California** - Do not apply unless parasite activity in the scale is at a low level or absent. (25 pints)

GRAPE: European Fruit Lecanium, Grape Leafhopper, Mealybug, Spider Mites - 1 pint. Do not apply within 3 days of harvest. Injury may occur to Ribier, Cardinal and Almeria varieties when applications are made after the clusters appear. (2¾ pints)

PEACH: Black Cherry Aphid, Black Peach Aphid, Green Peach Aphid, Japanese Beetle, Rusty Plum Aphid - ¾ pint. **Oriental Fruit Moth, Plum Curculio and Scales (European Fruit Lecanium, Terrapin, Cottony Peach)** - 1¼ pints. Do not apply within 7 days of harvest. (9 pints)

VEGETABLE AND FIELD CROPS

ALFALFA, CLOVER: Alfalfa Weevil Larvae, Aphids, Armyworms, Clover Leaf Weevil, Grasshoppers, Lygus Bugs, Pea Aphid, Potato Leafhopper, Spider Mites, Spittlebugs, Vetch Bruchid - 1¼ to 1½ pints. Use higher rate for Armyworm control. For hard to control insects, use up to 2 pints per acre. Apply to alfalfa in bloom only in the evening or early morning when bees are not working in the fields or are not hanging on the outside of hives. May be

7/12

applied on day of harvest or grazing up to 1½ pint rate; when using 2 pint rate, do not apply within 7 days of harvest or grazing. Do not apply during full bloom. (1½ pints)

ALFALFA, CLOVER, VETCH Grown for Seed (Not for use on VETCH in California):

Aphids, Leafhoppers, Lygus Bugs - 1 to 1¼ pints. Apply to plants in bloom only in the evening or early morning when bees are not working in the fields or are not hanging outside the hives. May be applied on the day of harvest.

BEAN: Aphids, Cucumber Beetle, Japanese Beetle, Mexican Bean Beetle, Nitidulid Beetles, Pea Leaf Weevil, Potato Leafhopper, Spider Mites - 1½ pints. **Lygus Bugs (on DRY BEANS in California)** - 1 to 1½ pints. Do not apply within 1 day of harvest. Do not graze or feed treated crop foliage to livestock.

BEET, GARDEN (Seed Crop): Lygus Bugs - 1¼ pints. Apply in seedball to hand seed stage. Repeat as needed. Do not apply within 7 days of harvest. (No limitation.)

BEET, TABLE: Aphids, Beet Armyworm, Blister Beetles, Flea Beetles - 2½ pints. Do not apply within 7 days of harvest. (2½ pints)

BERRIES (Blackberry, Blueberry, Boysenberry, Dewberry, Loganberry, Raspberry): Aphids, Japanese Beetle, Rose Leafhopper, Rose Scale, Two-Spotted Spider Mite, Sap Beetles - 2 pints. Do not apply within 1 day of harvest.

CELERY: Aphids, Leafhoppers, Spider Mites, Whiteflies - 1½ pints. Do not apply within 7 days of harvest. (1½ pints)

COLE CROPS (Broccoli, Brussels Sprout, Cabbage, Kale, Mustard, Turnip): Aphids, Cabbage Looper, Imported Cabbage Worm - 1½ to 2 pints. Use higher rate for loopers and worms. Do not apply within 3 days of harvest on Broccoli and Turnip, 7 days on others.

CORN: Aphids, Corn Rootworm Adults, Sap Beetles, Thrips, Young Grasshoppers - 1¼ pints. Do not apply within 5 days of harvest for forage use. **Caution NOTE:** Injury may occur in the whorl and silk stages with emulsifiable liquids. (1¼ pints)

COTTON: Brown Cotton Leafworm, Cotton Aphid, Cotton Leafworm, Cotton Leaf Perforator, Spider Mites, Leafhoppers, Lygus Bugs, Thrips, Whiteflies - ¾ to 1 pint. **Fall Armyworm, Boll Weevil, Grasshoppers, Garden Webworm, Lygus Bugs** - 1¼ to 2 pints. **Cotton Fleahopper** - ¾ to 1 pint. May be applied on the day of harvest. Consult local agricultural authorities for exact time of application.

CUCURBITS (Cucumber, Squash): Spider Mites, Squash Vine Borer, Thrips - 1¾ pints. Do not apply within 3 days of harvest on pumpkin. Do not apply on Cucumbers and Squash unless plants are dry or within 1 day of harvest. (Pumpkin 2 pints; others 1¾ pints.)

8/12

EGGPLANT: Aphids, Lace Bugs, Spider Mites - 1¼ to 2 pints. Use higher rate for Lace Bugs. Do not apply within 3 days of harvest.

GRASS AND CEREAL CROPS (Barley, Corn, Wheat, Barn Grass, Canary Grass, Fescue, Orchard Grass, Red Top, Timothy, Yellow Foxtail): Cereal Leaf Beetle - ¾ to 1¼ pints. Do not apply within 7 days of harvest or forage use on Barley and Wheat; 5 days on Corn. **Grass, Grass Hay Pastures and Range Land: Aphids, Leafhoppers** - 1 to 1¼ pints, repeat application as necessary. (Grasses 1¼ pints; Corn 1 pint; other cereals 1¼ pints)

HOPS: Aphids, Mites - 1¼ pints. Do not apply within 7 days of harvest. (1¼ pints)

HORSERADISH: Aphids, Diamondback Moth, Flea Beetles, Leafhoppers. Do not apply within 7 days of harvest. (2½ pints)

LETTUCE, ENDIVE: Aphids, Alfalfa Looper, Cabbage Looper, Leafhoppers, Mites - 2 pints. Do not apply within 14 days of harvest on leaf lettuce; 7 days on endive and head lettuce. (2 pints)

MINT: Aphids, Flea Beetle Adults, Spider Mites - 1 pint. Do not apply within 7 days of cutting. (1 pint)

ONION: Onion Maggot, Thrips - 2 pints. For Onion Maggot, apply when flies first appear and repeat every 4 days. Do not apply within 3 days of harvest on Green Onions. (2 pints)

PARSNIP: Aphids, Diamondback Moth, Flea Beetles, Leafhoppers - 2 pints. Do not apply within 7 days of harvest.

PEA: Aphids, Pea Weevils - 1¼ pints. Do not apply within 3 days of harvest. Do not graze or feed treated crop foliage to livestock.

PEPPER: Aphids, Pepper Maggot - 1½ pints. Do not apply within 3 days of harvest. (1½ pints)

POTATO: False Chinch Bug, Leafhoppers, Mealy Bugs - 1 pint. **Aphids, Blister Beetles** - 2½ pints.

RADISH: Aphids, Diamondback Moth - 2 pints. Do not apply within 7 days of harvest.

RICE: Leafminers - 1½ pints. Make first application when first rice blades appear on the surface of the water and repeat if necessary. **Stink Bugs** - 1 pint in a minimum of 2 gallons of water applied by aircraft during the early milk and dough stages of growing rice. Broadcast use only over intermittently flooded areas. Application may not be made around bodies of water where fish or shellfish are grown and/or harvested commercially. Do not apply within 7 days of

9/12

harvest. (1½ pints)

SALSIFY: Aphids, Diamondback Moth, Flea Beetles, Leafhoppers - 2 pints. Do not apply within 7 days of harvest.

SMALL GRAINS (Barley, Rye, Wheat): Armyworms, English Grain Aphid, Grasshoppers, Greenbugs - 1 to 1¼ pints. Do not apply within 7 days of harvest. (1¼ pints)

SPINACH: Aphids - 1½ pints. Do not apply within 7 days of harvest.

STRAWBERRY: Aphids (use at rate of 2 pints), **Field Crickets, Lygus Bugs, Potato Leafhopper, Spider Mites, Spittlebugs, Strawberry Leafroller, Strawberry Root Weevil, Thrips, Whiteflies** - 1½ to 2 pints. Do not apply within 3 days of harvest. (2 pints)

SWISS CHARD: Aphids, Blister Beetles - 2 pints. Do not apply within 7 days of harvest. (2 pints)

TOMATO: Aphids, Spider Mites - 1½ pints. **Drosophila Flies** - 2 pints in full coverage application to fruit and foliage. Do not apply within 1 day of harvest. At rates over 2 pints up to 3½ pints, do not apply within 5 days of harvest. **California only:** These higher rates may be applied up to 3 days of harvest. (3½ pints)

PASTURE AND RANGE LAND: Aphids, Grasshoppers, Leafhoppers - 1¼ pints. Or use 1¼ pints in 1 gallon diesel fuel oil by means of airplane or a turbine blower type sprayer. May be applied on day of harvest or grazing. (1¼ pints)

WASTELANDS, ROADSIDES, SOIL-BANK LANDS (Not Grazed): Grasshoppers - 1 to 2 pints in 1 gallon fuel oil by means of an airplane or a turbine blower type sprayer.

STORAGE PEST PROTECTION

For protection against: **Cereal Leaf Beetle, Confused Flour Beetle, Indian Meal Moth, Flat Grain Beetle, Granary Weevil, Lesser Grain Borer, Red Flour Beetle, Rice Weevil, Rusty Grain Beetle, Sawtoothed Grain Beetle.**

GRAINS (Barley, Corn, Oats, Rye, Wheat) - RESIDUAL SPRAY - BEFORE STORING: (For walls, floors and machinery in grain elevators, in treating truck beds, boxcars, and ships' holds before loading grain): Use 5 pints per 25 gallons of water. Make thorough application. Before applying spray, clean elevators, boxcars, truck beds, etc., thoroughly. Remove and burn all sweepings and debris.

ORNAMENTALS

Dosages given are for use in 100 gallons of dilute spray.

10/12

Aphids, European Pine Shoot Moth, Four-Lined Leaf Bug, Japanese Beetle Adult, Mealybugs, Millipedes, Potato Leafhopper, Rosa Leafhopper, Scurfy Scale, Spider Mites, Springtails, Scwbugs, Tarrished Plant Bug, Thrips, Whiteflies, Euonymus Scale, Lace Bugs, Oyster Shell Scale - 1 pint.

Azalea Scale, Bagworm, Birch Leafminer, Boxwood Leafminer, Fletcher Scale, Florida-Red Scale, Juniper Scale, Magnolia Scale, Oak Kermes, Pine Leaf Scale, Tent Caterpillar - 1 1/4 pints.

Black Scale Crawlers, Monterey Pine Scale - 1 3/5 pints.

Pine Needle Scale, Wax Scale - 2 pints.

For **Oyster Shell, Fletcher, Juniper, Oak Kermes and Pine Needle Scales**, apply when scale crawlers have settled on foliage.

CAUTION NOTE: Avoid use on certain ferns including Boston, Maidenhair and Pteris, as well as some species of Crassula and Canaerti Juniper.

FLY CONTROL

For use around the outside of buildings which house domestic animals, around the outside of homes, around yards, and the outside of meat and food processing plants.

Straight Atrapa 8E Sprays

<u>Amount of Spray</u>	<u>Amount of Atrapa 8E</u>
2 1/2 gal.	3/4 cup
12 gal.	1 1/4 pint
100 gal.	1 1/4 gal.

Atrapa 8E Bait Sprays

<u>Amount of Bait Spray</u>	<u>Amount of Atrapa 8E</u>	<u>Sugar (or)</u>	<u>Unsulfured Molasses or Corn Syrup</u>
2 1/2 gal.	3/4 cup	1 cup	1 cup
12 gal.	1 1/4 pt.	2 1/2 lbs.	1 qt.
100 gal.	1 1/4 gal.	20 lbs.	2 gal.

Apply spray at rate of 1 gallon per 1,000 square feet on painted surfaces and 2 gallons per 1,000 square feet on unpainted surfaces where flies alight or congregate. In most cases, adding molasses or sugar to the spray prolongs the insecticidal activity and serves as a fly attractant.

CAUTION NOTE: Avoid contamination of milk, milk equipment and water. Avoid contamination of feed and food products; also drinking fountains and feed troughs.

DROSOPHILA FLIES and DRIED FRUIT BEETLE: On or around cull fruit and vegetable dumps. Mix 7 1/2 pints in 100 gallons of water and apply as a drench, using 8 to 10 gallons of spray per 100 square feet (for best results, dumps should not be over 18 inches deep). Do not feed treated fruit and vegetables.

11/12

MOSQUITO CONTROL

MOSQUITOES, FLIES AND SMALL FLYING INSECTS: (Broadcast use only over intermittently flooded areas. Application may not be made around bodies of water where fish or shellfish are grown and/or harvested commercially.) For use by trained personnel as a 2% to 5% fog, aerosol or space spray. To make a 2% solution, dilute 1 part Atrapa 8E in 45 parts of water, fuel oil or diesel oil. When using kerosene-type solvent as carrier, dilute 1 part Atrapa 8E in 45 parts of a mixture consisting of 4 parts kerosene-type solvent and 1 part aromatic hydrocarbon-type solvent. For a 5% solution, dilute 1 part Atrapa 8E in 18 parts of mixture using similar solvents.

MOSQUITO LARVAE IN STANDING WATER: (Broadcast use only over intermittently flooded areas. Application may not be made around bodies of water where fish or shellfish are grown and/or harvested commercially.) Apply at the rate of 8 fluid ounces (approximately ½ pint) per acre. Mix in sufficient water or oil to obtain even coverage when applied by air or ground. Repeat applications as necessary. Avoid applying oil-based formulations to valuable ornamental plants as injury may occur. Keep out of any fish bearing waters.

ADULT MOSQUITO AND FLY CONTROL

Apply 2.4 to 4.8 fluid ounces per acre for control of Adult Mosquitoes and 7.2 to 9.6 fluid ounces per acre for control of Adult Flies and Mosquitoes. Repeat as necessary. Can be applied to Alfalfa, Clover, Pasture and Range Grass, Grass and Grass Hay, Grain Crops (Barley, Corn, Oats, Rye and Wheat), Beans, Rice and Nonagricultural Lands. Do not apply within 7 days of harvest on Grain Crops (Barley, Oats, Rye and Wheat) and Rice; 5 days of harvest on Corn; 1 day of harvest on Beans. No time limitation on Alfalfa, Clover, Pasture and Range Grass, Grass and Grass Hay. Do not apply to Alfalfa and Clover in bloom. Do not use on Seed Alfalfa. Treat shrubbery and vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, ponds and shorelines may be treated. (Broadcast use only over intermittently flooded areas. Application may not be made around bodies of water where fish or shellfish are grown and/or harvested commercially.)

Mist blowers and boom sprayers utilizing a controlled air flow to facilitate particle size and spray deposition may be used at a vehicle speed of 4 to 10 mph.

Mist blowers with a pump capable of producing up to 40 psi and blower speeds of 2600 rpm are satisfactory. Use flat fan nozzles, 8001 to 8002, placed 30° into air blast or rotary atomizers into the air blast that produce an efficient spray particle with a mass medium diameter of 40 to 100 microns. Swath widths should not exceed 30 feet, and applications should not be made when winds exceed 5 mph.

Repeat applications should be made as necessary unless otherwise specified.

12/12

WARRANTY STATEMENT

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. The exclusive remedy of any buyer of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall not exceed the purchase price paid for this product or, at Griffin Corporation's election, the replacement of this product. **GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

Atrapa is a registered trademark of Griffin L.L.C.
Griffin and Design are a registered trademark of Griffin Corporation.

[Based on the EPA stamped accepted label (with comments) dated February 10, 1999]