

ATOMIC



WARFARIN

U.S.D.A. Registration No. 4789-1
CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTION ON BACK PANEL

CONCENTRATED

Contents Makes 10 lbs.

RAT
AND MOUSE
KILLER BAIT

Active Ingredients
Inert Ingredients

**"ATOMIC DOES NOT
DEVELOP BAIT SHYNESS"**

ACCEPTED
6-21-68




U.S.D.A. Registration No. 4789-1
CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER SIDE OF BACK PANEL


CONCENTRATED
Contents Makes 10 lbs.



RAT
AND MOUSE



KILLER BAIT



**ATOMS DOES NOT
DEVELOP BAIT SHYNESS**

DIRECTIONS

Mix thoroughly one part of ATOMIC WARFARIN with 19 parts, by weight, of suitable bait material such as fresh corn meal, rolled oats, poultry mash, etc. Entire contents may be mixed at once by adding 9 $\frac{1}{2}$ lbs. of any mixed ground feed.

for all rat and mouse infestations—

DIRECTIONS FOR USE

GENERAL DIRECTIONS: Place baits in location where rats and mice feed, water or travel. Bait replacements along walls, in corners, and in concealed places where rats and mice can feed unobserved are desirable. Contaminated or spoiled bait should be replaced promptly. Where a continuous source of infestation is present, permanent bait stations should be established and replenished as needed.

RATS: Place 4 to 6 oz. of bait in dry containers where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 10 days and continue baiting until all signs of feeding have stopped.

MICE: Place tablespoon amounts ($\frac{1}{4}$ to $\frac{1}{2}$ oz.) where mice or signs of mice have been seen. Bait stations of 12 to 15 foot intervals are recommended. An uninterrupted supply of bait should be maintained for at least 15 days.

Baits should be placed in locations not accessible to children, pets and domestic animals, or in tamper proof bait boxes.


CAUTION

Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

MOON CHEMICAL PRODUCTS

Martville

New York



With Comments