

CHLORMADANE

FOR CONTROL OF TICKS, LICE, FLEAS, AND SARCOPTIC AND
DEMODECTIC MANGE OF DOGS

DIRECTIONS

For Fleas and Lice: Add 2 ounces (4 tablespoonfuls) Chlormadane to 1 gallon of water. Animal should first be bathed with soap and water, rinsed and toweled. Apply the diluted Chlormadane solution with sponge or cloth or by dipping and allow the animal to dry without toweling. To prevent reinfestation, repeat application every 2 to 3 weeks.

For Ticks: Add 4 ounces (8 tablespoonfuls) Chlormadane to 1 gallon of water and follow above directions.

Since diagnosis of demodectic mange is often difficult, and treatment may involve other factors such as diet, treatment should be done by or under direction of a veterinarian.

ACTIVE INGREDIENTS:

Technical Chlordane** -----	4.0% w/w
Gamma isomer of benzene hexachloride from Lindane --	1.0% w/w
Malathion (O,O-dimethyl dithiophosphate) -----	1.0% w/w
Petroleum solvent -----	58.5% w/w

INERT INGREDIENTS ----- 35.5% w/w

Total ----- 100.0% w/w

**Refined (clarified grade) equivalent to 2.4% Octachloro-4, 7-methanotetrahydroindane and 1.6% related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN. FOR EXTERNAL USE ONLY.
Use only the diluted solution as recommended. Harmful if swallowed.
Avoid inhalation or contamination of feed. Repeated or prolonged contact with skin can cause toxic symptoms. Wash thoroughly after using. DO NOT use on cats or nursing puppies.

MADE IN U.S.A.

U.S.D.A. REG. NO. 4812-

CENTER PANEL:

CHLORMADANE

Net Contents _____

FOR CONTROL OF FLEAS, LICE AND TICKS

CAUTION: Keep Out of Reach of Children.

Manufactured by:
Eastern Laboratories, Inc.
1483 Washington Avenue
Vineland, New Jersey 08360