

CODE 820.75

PYRENONE® Multi-Purpose Insecticide

Contains PYRENONE®

Professional Strength - Low Dosage Required

For use in Schools, Hospitals, Institutions, Supermarkets, Dairies, Hotels, Restaurants, Drive-ins, Motels, Drug Stores, Food Plants, Milk Rooms and Animal Quarters

Provides Rapid Control of Flies, Mosquitoes, Flying Moths (Millers), Gnats, Wasps and accessible stages of Roaches, Ants, Water Bugs, Silverfish, Flour Beetles, Weevils, Grain Mites and Spiders

KILLS AND REPELS Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats

Controls Garden Insects on Ornamental and Vegetable Plants

Acceptable for Use to Day of Harvest on Edible Plants

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
Petroleum Hydrocarbons	4.3%
INERT INGREDIENTS:	90.2%
	100.0%

*Equivalent to 4% (butylcarbityl)(6-propylpiperonyl) ether and 1% related compounds

Keep Out Of Reach Of Children

CAUTION

See side panel for additional precautionary statements.

PYRENONE - Registered trademark of Fairfield American Corporation

DOES NOT CONTAIN CHLOROFLUOROCARBONS

EPA Reg. No. 4816-525
EPA Est. No. 279-NY-1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid breathing of vapor or contact with eyes or skin. Vacate premises after treatment and ventilate before reoccupying.

Avoid contamination of feed and foodstuffs. When using this product in food processing plants, food should be removed or covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

Do not apply over uncovered milk or food products. Any exposed utensils or food handling surfaces should be thoroughly cleaned before using.

Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

Do not use in areas where fish or reptiles may be exposed to the spray.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

DISPOSAL: Wrap container and put in trash collection.

SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.



4816-525

12-3-78

10/2

**FAIRFIELD
AMERICAN
CORPORATION**

3932 Salt Road
Medina, New York 14103

Flies, Mosquitoes, Gnats, Wasps and Flying Moths (Millers): Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat as necessary.

Roaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes and Sow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Exposed Stages of Indian Meal Moths and Angoumois Grain Moths: Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Exposed Stages of Flour Beetles, Weevils and Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

To Treat Animals: To protect beef and dairy cattle from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats, spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

In Animal Quarters and Milk Rooms: For rapid control of house flies, mosquitoes, gnats, wasps and small flying moths, close windows and doors and spray at the rate of 1 second per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

For Use in Gardens: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

USE ON	TO CONTROL
Roses	Aphids, Rose Chafers, Japanese Beetles, Rose Slugs, Tent Caterpillars, Scales
Asters	Leafhoppers, Tarnish Plant Bugs, Lace Bugs, Thrips (exposed), Mealy Bugs
Dahlias	Aphids, Red Spider Mites, Plant Bugs
African Violets	Broad Mites, Citrus Mealybugs
Azaleas	Azalea Lace Bugs, White Flies, Leaf Miners
Euonymus	Euonymus Scales, Aphids, Thrips (exposed)
Hydrangeas	Aphids, Red Spider Mites, Hydrangea Leaf-tiers
Junipers	Juniper Scales, Red Spider Mites, Bagworms
Lilacs	Leaf Miners, Leaf Rollers, Scales
Tomatoes	Flea Beetles, Hornworms, Stink Bugs, Colorado Potato Beetles, Green Peach Aphids
Peppers, Eggplants, Potatoes	Colorado Potato Beetles, Leafhoppers, Blister Beetles, Flea Beetles, Green Peach Aphids
Cucumbers, Squash, Melons	Striped & Spotted Cucumber Beetles
Beans	Mexican Bean Beetles, White Flies, 12-Spotted Beetles
Sweet Corn	Japanese Beetles

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

2092