

CODE 723.30

PYRENE

EPA Reg. No. 411-100
EPA Est., 279-NY-1

For Outdoor Fogging Application in Residential Areas, Parks, Grounds, Golf Courses, Municipalities and Schools, and Restaurants

Kills Mosquitoes, Gnats, Flies and Bees

As a Contact Spray indoors in Outdoors kills Ants, Fleas, Crickets, and Box Elder Bugs. Use as a Spot Spray for Fleas avoiding any Contact with Food or Food Products

ACTIVE INGREDIENTS

- Pyrethrins
- *Piperonyl Butoxide, Technical
- **Malathion
- Aromatic Petroleum Derivative Solvent
- Petroleum Distillate

*Equivalent to 0.100% (butylcarbitol) 6-propylperoxy ether or to 0.025% related compounds

**0,0-Dimethyl dithiophosphate of diethylmercaptosulfonate

PYRENONE Fairfield American trademark

CAUTION
KEEP OUT OF REACH OF CHILDREN.

CAUTION

Harmful if swallowed. Avoid breathing of fumes. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feet and foodstuffs

Do not use in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or served. Do not use in serving areas while food is exposed

This product is toxic to fish. Do not apply when weather conditions favor drift from areas treated. Keep out of ponds, streams, lakes, and not contaminate lakes, streams or ponds by cleaning or equipment or disposal of wastes

Do not reuse empty container. Do not use perforated containers and burying in a safe place or return to drum recycler

For Outdoor Fogging Application in Residential Areas, Parks, Grounds, Golf Courses, Municipalities and Schools, and Restaurants

As a Contact Spray indoors in Outdoors kills Ants, Fleas, Crickets, and Box Elder Bugs. Use as a Spot Spray for Fleas avoiding any Contact with Food or Food Products

For outdoor fogging application, the above application rate should be used when traveling at a speed of 5 m.p.h. at a 400 foot nozzle height. Treat with insecticide dispersed at a rate of from 1 to 2 gallons per hour.

For outdoor contact spray application, the above application rate should be used when traveling at a speed of 5 m.p.h. at a 400 foot nozzle height. Treat with insecticide dispersed at a rate of from 1 to 2 gallons per hour.

For indoor contact spray application, the above application rate should be used when traveling at a speed of 5 m.p.h. at a 400 foot nozzle height. Treat with insecticide dispersed at a rate of from 1 to 2 gallons per hour.

For indoor contact spray application, the above application rate should be used when traveling at a speed of 5 m.p.h. at a 400 foot nozzle height. Treat with insecticide dispersed at a rate of from 1 to 2 gallons per hour.

NET CONTENTS	_____	GALLONS
NET CONTENTS	_____	LITERS
NET CONTENTS	_____	POUNDS
NET CONTENTS	_____	KILOGRAMS

ACCOUNTED

BATCH _____

DATE 02 1977

FUNDED AND REGISTERED UNDER NO. _____



FAIRFIELD AMERICAN CORPORATION
100 N. AGARA ST.
MIDDLEPORT, N.Y. 14105