PLEASE NOTE

This image contains more than one label approved for this product on this date.

.1

2

4	4816	- 331	61	1/1978	····· }	- por	c 181
					C C C C C C C C C C C C C C C C C C C		
	NET CONTENTS 1 QUART NET CONTENTS 2.08 POUNDS ATE	<pre>IMDOR USE This spray may be used in food processing plants. homes, restaurants, hospitals, torres, hose is motels, schools and other public health buildings, motels, schools and other public health buildings, had biries, barrs, cannerles, brewerles and food had biries, barrs, cannerles, brewerles and food had biries, barrs, samil flying moths flies hosquitoes Dilute 2 tableapoons or 0.6 sunces of concentrate per gallon of water or oil. Ramove pet, birds and cover fish aquatia before spraying.</pre>	A control of a start of the start and start of the start and a start of the start and the start of the start	Cheese Stippers' Silverfish Cheese Altess' Spiders Spiders Dilute at the rate of 2 oundes of condentruction per gallon of water of oil. Mapty of a four face strain in ground matcher abuited to of diver a coarte spary at the opticular of diver a coarte feet. In Porcularly aget wallow of States and crucks where inserts inder. Spea ond runs. India Lastebander, herehold, Sod		· · · · · · · · · · · · · · · · · · ·	
	SPRAY CONCENTRATE	COUND APLICATION adult Mosquices Gnats Small Flying Moths Small Flying Moths Small Flying Hoted Application Faces: <u>Jurion and application Faces:</u> <u>Jurion atter</u> <u>Bliuted Area to be</u> te <u>outor</u> <u>Jil.or.</u> 2000 sq.ft		statistic from the state in the state may be hour, dis- persing 125 gallons of sprey per hour. dis- persing 125 gallons of sprey per hour. dis- To control <u>morquito larvae</u> : Dilute at the rate of 7 ounces per 50 gallons of oil and apply at the rate of 20 to 25 gallons per acte. Apply as a fog or fine mist uniformly weter where larvae may break of on ot acceed this weter where larvae may break of on ot acceed this	dosage or fish kill may result. Alactarf AppLication: FixED VING AlactarFT To control: Adult mosquitoes Dilute 1 to 2 ounces of concentrate in 4 gallons of water per acre.		
	C C E MGSOUITO S	Muture the Frankrich Tresection. Frankriche, dra Rossergeschurgenn Insterents an archarde, dra Rossergeschurgenn Insterents großserstein in food and the archard article and the metal and the archard article articles are articles and the archard articles are before using. When using the product in these soon perfection. ENVIRONMENTAL FORMANDS	This product is toxic to fish, keep out of lakes streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wates. Ap- ply this product only as specified on this label Co not apply outdoors when weather conditions favor drift from areas treated.	It is a violation of Federal law to use this pro- duct in a manner inconsistent with its labeling. STORAGE AND OISPOSAL Bo not reuse empty container. Wrap container and put in trash collection.	This is a highly concentrated product designed to be diluced before use in either water or oil. The resulting spray should be dispersed through good fogging or spraying equipment.		
	CODE 723.70 EPA Reg. No. 4816-331 EPA Est. No. 279-NY-1 JUN	A highly concentrated Pyrenome Insect Lighture the P Dilute in mater or oil before use cas activities for outdoor or Indoor application as registry and Spray KTIVE INGREDIENTS. Pretrolement Distributes Perrolement Distributes Nettilizate (continue continue continu	IOU.CO Equivalent to 48 (burytearbityt)(6-propytaip- eronyt) sther and 12. related compounds CONTON ALACH OF CHILDRIN. CAUTION Harnful if swallowed. See side panel for additional precautionary statements.	PYRENONE - Registered trademark of Fairfield American Corporation BATCH	72370 10 278		

an an Star Star An Star Star Star An Star Star Star

LABEL

,

÷.

NEXT

CODE 723.70

MOSQUITO SPRAY CONCENTRATE

A highly concentrated Pyrenone[®] Insecticide

Dilute in water or oil before use.

For outdoor or indoor application as a Fog or Spray

ACTIVE INGREDIENTS:

Pyrethrins	6.0%
*Piperonyl Butoxide, Technical	
Petroleum Distillate	20.0%
INERT INGREDIENTS:	14.0%
	100.0%

*Equivalent to 48% (butylcarbityl)(6propylpiperonyl) ether and 12% related compounds

Keep Out Of Reach Of Children

CAUTION

Harmful if swallowed. See side panel for additional precautionary statements.

EPA Reg. No. 4816-331 EPA Est. No. 279-NY-1

PYRENONE - Registered trademark of Fairfield American Corp.

NET CONTENTS NET CONTENTS NET CONTENTS NET CONTENTS	GALLONS LITERS POUNDS KILOS	
Buyer assumes all risks of or handling of this materi. strict accordance with dir herewith. BATCH	use, störage al not n	*
		CAR 32 E adin

·
and the second
and the second sec
and a second sec
·
and the second sec
and a second sec











Apply as a fog or fine mist. Treat entire area with particular attention to shrubbery and vegetation where insects may rest, also low or marshy areas, shore lines and stagnant pools or ponds. Repeat as necessary, usually once or twice a week.

With mobile or truck mounted equipment, the above application rate will result, assuming a 400 foot swath width, traveling at 5 miles per hour, dispensing 125 gallons of spray per hour.

4816

١

 $\hat{\boldsymbol{\omega}}$

11/19-2

ιÚ

To control: Mosquito Larvae

Dilute at the rate of 7 ounces per 50 gallons of nil and apply at the rate of 20 to 25 gallons per acre. Apply as a fog or fine mist uniformly over wet lands, swamps, marshes or bodies or water where larvae may breed. Do not exceed this dosage or fish kill may result.

AIRCRAFT APPLICATION: FIXED WING AIRCRAFT To control: Adult Mosquitoes

Dilute 1 to 2 ounces of concentrate in 4 . gallons of water per acre. INDOOR USE:

This spray may be used in food processing plants, homes, restaurants, hospitals, stores, hotels, motels, schools and other public health buildings, in dairies, barns, canneries, breweries, and food handling establishments.

To control: Gnats Small Flying Moths Flies Mosquitees

Dilute 2 tablespoons or 0.6 ounces of concentrate per gallon of water or oil. Remove pets, birds and cover fish aquaria before spraying. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of room, thoroughly filling the area with mist. Do not remain in treated areas. Ventilate the area after treatment is completed. Repeat treatment when reinfestation occurs.

To control accessible stages of: Ants Cheese Mites Cockroaches Grain Beetles Crickets Granary Weevils Confused Flour Beetles Rice Weevils Cadelles Silverfish Cheese Skippers Spiders

Rust-red Fiour Beetles Saw-toothed Grain Beetles Dilute at the rate of 2 ounces of concentrate per gallon of water or oil. Apply as a surface spray in a good machine adjusted to deliver a coarse spray at the rate of 1 gallon to 750 square feet. Thoroughly spray walls, floors, shelves, pallets with particular attention to cracks and crevices where insects hide. Spray ant runs, trails, baseboards, thresholds and other places of entry and treat nests liberally. Repeat when needed.

ACCEPTED JUN 1 1978 Under the Peneral Insecticity. Particulates and Hadron Fridde Act. as amended, for the penetodia

4616 - 33

et hin R

2

•

out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

> Do not apply outdoors when weather conditions favor drift from areas treated.

ENVIRONMENTAL HAZARDS This product is toxic to fish. Keep

not in operation.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is

......

242

DIRECTIONS FOR USE It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STURAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be dispos ed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Reseal container and offer for reconditioning or triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved lundfill or bury in a safe place. Consult Federal, state or local disposal authurities for approved alternative procedures.

Use the following dilution and applica-

bilut		Use Rate		
oncentrate	Water or Oil	Diluted Spray	Area to be Covered	
3 this.	1 quert .	3 fl. oz.	2000 sq.ft.	
51 fl.oz.	1 gallon	12 fl. oz.	8000 sq. ft.	
2 gallons	50 gallons	1 gallon	1 acre	

72370 278