PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

Do not apply outdoors when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage of disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or Chemically processed should be dispoeset of in a landfill approved for a pesticides on buried in a safe place away from Water supplies. Reseal container and offer for reconditioning or triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place. Consult Federal, state or local disposal authorities for approved alternative procedures.

A highly concentrated Pyrenone® Insecticide

MOSQUITO SPRAY CONCENTRATE

Dilute in water or oil before use.

For outdoor or indoor application as a Fog or Spray

ACTIVE INGREDIENTS.

		•						
Pyrethrins	<i>.</i>							6.0%
"Piperonyl	Butoxi	de.	Tec	chr	١i٥	a	1	60.02
Petroleum	Distil	late						20.0%
INERT INGREC	HENTS:							14.0%
								100.02

*Equivalent to 48% (butylcarbity))(6propylpiperonyl) ether and 12% related compounds

EPA Reg. No. 4816-331 EPA Est. No. 279-NY-1

PYRENONE - Registered trademark of Fairfield American Corp.

Keep Out Of Reach Of Children

CAUTION

Harmful if swallowed. See side panel for additional precautionary statements.

NET CONTENTS	GALLONS
NET CONTENTS	LITERS
NET CONTENTS	POUNDS
NET CONTENTS	KILOS

BATCH

OUTDOOR GROUND APPLICATION:

To control: Adult Mosquitoes Small Flying Moths

Use the following dilution and application rates:

Dilution		Use Rate				
oncentrate	Water or 011	Diluted Spray	Area to be Covered			
3 tbls.	1 quart	3 fl. oz.	2000 sq.ft.			
5} fl.oz.	l gallon	12 ft. az.	8000 54. ft.			
2 gallons	50 gallons	1 galton	acre			

Apply as a fog or fine mist. Treat entire area with particular attention to shrubbery and vegetation where insects may rest, also low or marshy areas, shore lines and stagnant pools or ponds. Repeat as necessary, usually once or twice a week.

With mobile or truck mounted equipment, the above application rate will result. assuming a 400 foot swath width, traveling at 5 miles per hour, dispensing 125 gallons of spray per hour.


To control: Mosquito Larvae

Dilute at the rate of 7 ounces per 50 gallons of oil and apply at the rate of 20 to 25 gallons per acre. Apply as a f or fine mist uniformly over wet lands, swamps, marshes or bodies or water where larvae may breed. Do not exceed this dosage or fish kill may result.

AIRCRAFT APPLICATION: FIXED WING AIRCRAFT

To control: Adult Mosquitoes

Dilute 1 to 2 ounces of concentrate in 4 gallons of water per acre.







FAIRFIELD AMERICAN CORPORATION

3932 Salt Road Medina, New York 14103



INDOOR USE:

This spray may be used in food processing plants, homes, restaurants, hospitals, stores, hotels, motels, schools and other public health buildings, in dairies, barns, canneries, breweries, and food handling establishments.

To control: Gnats Small Flying Moths
Flies Mosquitoes

Dilute 2 tablespoons or 0.6 ounces of concentrate per gallon of water or oil. Remover pets, birds and cover fish aquamica before spraying. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of room, thoroughly filling the area with mist. Do not remain in treated areas. Ventilate the area after treatment is completed. Repeat treatment when reinfestation occurs.

To control accessible stages of:

Ants Cheese Mites
Cockroaches Grain Beetles
Crickets Granary Weevils
Confused Flour Beetles Rice Weevils
Cadelles Silverfish
Cheese Skippers Spiders
Rust-red Flour Beetles
Saw-toothed Grain Beetles

Dilute at the rate of 2 ounces of concentrate per gallon of water or oil. Apply as a surface spray in a good machine adjusted to deliver a coarse spray at the rate of 1 gallon to 750 square feet. Thoroughly spray walls, floors, shelves, pallets with particular attention to cracks and crevices where insects hide. Spray ant runs, trails, baseboards, thresholds and other places of entry and treat nests liberally. Repeat when needed.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

2