

Rose and Floral Spray

Residual

A Special Blend of Insecticide, Fungicide, and Miticide

EPA Reg. No. 4816-348 EPA Est. No. 279-NY-1

Active Ingredients

Pyrethrins	0.025%
*Piperonyl Butoxide, Technical	0.256%
Rotenone	0.128%
Other Cube ¹ Extractives	0.237%
Folpet: N-(trichloromethylthio) phthalimide	0.700%
Carbaryl: 1-naphthyl N-methylcarbamate	1.000%
Petroleum Distillate	0.025%
Inert Ingredients	97.629%
	<hr/>
	100.000%

*Equivalent to 0.205% (butylcarbaryl) (6-propylpiperonyl) ether and to 0.051% related compounds

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or inhaled. Avoid breathing spray or contact with eyes or skin. Wash thoroughly after use. Avoid contamination of feed or foodstuffs.

_____ **Pounds Net Weight**

_____ **Kilograms Net Weight**


Do not spray closer than 18 inches from plants. Do not use near fish-bowls or aquaria.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

Terms of Sale or Use On purchase or use of this product buyer and user agree to the following conditions

Warranty FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **Except as so warranted the product is sold as is. FMC makes no other warranty express or implied.**



FMC Corporation
A.C.D. Industrial Sales Department
Middletown, New York 14105

Directions for use

Hold container at least 18" from foliage, spray lightly in one-second bursts, spraying both top and underside of foliage. Injury to tender foliage may occur if plants are wet when treated or in the presence of high humidity.

For control of the insects listed with the ornamentals below, treat when infestation occurs. Properly treated, foliage will be protected from the specified insects for about one week. See other specific directions for listed ornamentals.

Ornamentals	Insects	Diseases	Other Specific Directions
African Violets	Mites Mealybugs	Powdery mildew Gray mold blight	
Asters	Thrip (exposed) Mealy bug Leafrollers Aphids Leafhoppers Plant bug Lace bug Blister beetle	Powdery mildew	Powdery mildew: begin before disease appears. Repeat at 7 to 10 day intervals, more frequently during rains or high humidity.
Azaleas	Lace bug Thrip (exposed) Mite White fly Scale Mealy bug		
Begonias	Mealy bug Thrip (exposed) Mites White fly Scale Orange tortrix		
Carnations	Mites Thrip (exposed) Aphid Mealy bug	Alternaria leaf spot Rust	Alternaria leaf spot, and rust: Begin two to three weeks after planting, repeat at 7 to 10 day intervals as necessary.
Chrysanthemums	Aphid Leafhoppers Thrip (exposed) Mites Plant bug Mealy bug White fly Scale Orange tortrix	Powdery mildew Alternaria leaf spot	Powdery mildew, and Alternaria leaf spot: Begin immediately after planting, repeat at 7 to 10 day intervals, more frequently during rains or high humidity.
Gladioli	Aphid Leafhopper Leaf spot Mite Thrip (exposed)		
Impatiens	White fly Scale		

Ornamentals	Insects	Diseases	Other Specific Directions
Hydrangeas	Aphid Scale Plant bugs Mites	Didymelasma leafspot	Didymelasma leafspot: Begin when plants emerge, repeat at 7 to 10 day intervals.
Lilies	Thrip (exposed) Leafhopper Japanese beetle Scale Orange tortrix	Powdery mildew	Powdery mildew: apply before disease appears or at first signs, repeat as necessary.
Marigolds	Japanese beetle Plant bug Leafhoppers Mites	Alternaria leafspot	Alternaria leafspot: apply at 3 to 7 day intervals during periods of rapid growing or high humidity.
Phlox	White fly Mealy bug Mites Thrip (exposed) Aphids	Powdery mildew	Powdery mildew: begin before disease appears. Repeat at 7 to 10 day intervals or more frequently during rains or high humidity.
Rhododendrons	Lace bug Mealy bug Aphid Scale Thrip (exposed) White fly		
Roses	Aphid Leafhopper Rose chafe Japanese beetle Rose slug June bug Leafroller Fall webworm Mites Plant bug Scale White fly Mealy bug	Black spot Powdery mildew	Black spot, powdery mildew: begin when first leaves unfold, repeat treatment at 7 to 10 day intervals through the growing season.
Rubber plant	Lace bug Scale Mealy bug Thrip (exposed)		
Sand daisy	Mite Plant bug Aphid	Alternaria leaf spot Powdery mildew Rust	Alternaria leaf spot, powdery mildew, and rust: Begin when plants emerge, repeat at 7 to 10 day intervals.
Wendy's plant	Scale Mealy bug Mite		
Zinnias	Japanese beetle Leafhopper Scale Mite Thrip (exposed) Plant bug Mealy bug White fly Scale Orange tortrix	Alternaria leaf spot	Alternaria leaf spot: Begin when plants emerge, repeat at 7 to 10 day intervals.