

Reg # 4816-348 PM-12

181

CODE 824.10

# ROSE AND FLORAL SPRAY

## RESIDUAL

EPA Reg. No. 4816-348 EPA Est. No.

A Special Blend of Insecticide, Fungicide and Miticide

### ACTIVE INGREDIENTS

Pyrethrins .....	0.025%
* Piperonyl Butoxide, Technical .....	0.256%
Rotenone .....	0.128%
Other Associated Resins .....	0.237%
Folpet* N-Trichloromethylthio phthalimide .....	0.700%
Carbaryl (1-Naphthyl N-Methylcarbamate) .....	1.000%
**INERT INGREDIENTS .....	97.654%
	100.000%

\*Equivalent to 0.205% (butylcarbaryl)(6-propyl-piperonyl) ether and 0.051% related compounds

\*\*Contains Petroleum Distillate

**KEEP OUT OF REACH OF CHILDREN.**  
**CAUTION**  
 See rear panel for use cautions.

This product contains no chlorofluorocarbons

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hold container at least 18 inches from foliage, spray lightly in one second bursts, spraying both top and underside of foliage. Injury to tender foliage may occur if plants are wet when treated or in the presence of high humidity. Do not spray blooming African Violets or Wax Begonias.

For control of the insects listed with the ornamentals below, treat when infestation occurs. Properly treated, foliage will be protected from the specified insects for about one week. See other specific directions for listed ornamentals.

ORNAMENTALS	INSECTS	DISEASES	OTHER SPECIFIC DIRECTIONS
African Violets	Mites	Powdery Mildew Gray Mold Blight	
Azaleas	Thrips (Exposed) Mealy Bugs Leafrollers Aphids Leafhoppers Plant Bugs Lace Bugs Blister Beetles	Powdery Mildew	Powdery Mildew: Begin before disease appears. Repeat at 7 to 10 day intervals, more frequently during rains or high humidity.
Asclepias	Lace Bugs Thrips (Exposed) Mites White Flies Scale Leafy Bug		
Begonias	Mealy Bugs Thrips (Exposed) Mites White Flies Scale Orange Tortrix		
Carnations	Mites Thrips (Exposed) Aphids	Alternaria Leaf-spot Rust	Alternaria Leafspot and Rust: Begin two (2) weeks after planting, repeat at 7 to 10 day intervals as necessary
Chrysanthemums	Aphids Leafhoppers Thrips (Exposed) Mites Lace Bugs Plant Bugs Mealy Bugs White Flies Scale Blister Beetles	Powdery Mildew Septoria Leafspot	Powdery Mildew, Septoria Leafspot: Begin immediately after planting, repeat at 7 to 10 day intervals, more frequently during periods of high humidity.
Dahlias	Aphids Plant Bugs Leafhoppers Mites Thrips (Exposed)		
Geraniums	White Flies Mealy Bugs Mites Leafrollers Webworms		
Iris	Aphids Scale Plant Bugs Mites	Didymellina Leafspot	Didymellina Leafspot: Begin when plants emerge, repeat at 7 to 10 day intervals
Lilac	Thrips (Exposed) Leafrollers Japanese Beetles Scale Orange Tortrix	Powdery Mildew	Powdery Mildew: Apply before disease appears or at first signs, repeat as necessary
Marigolds	Japanese Beetles Plant Bugs Leafhoppers Mites	Alternaria Leaf-spot	Alternaria Leafspot: Apply at 3 to 7 day intervals during periods of rapid growing or high humidity

ACCEPTED

MAR 05 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under Reg. No. 4816-348

82410 1184 NET CONTENTS:



CODE 824.10

FAIRFIELD

AMERICAN CORPORATION

239 WILSON AVENUE  
NEWARK, N.J. 07105