- Highly Active Synthetic Pyrethroid
- Effective in Controlling a Wide Variety of Insect Pests in the Household and on Ornamental Garden Plants.

ACTIVE INGREDIENT:

*Resmethrin	0.25%
INERT INGREDIENTS:	99.75%
	100.00%

*Cis/trans isomers ratio: Max 30% (±) cis and min 70% (±) trans. Synthrin - Registered Trademark of AgrEvo Environmental Health, Inc.

EPA Reg. No. 4816-382

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious. do not give anything by mouth and do not induce vomiting.

If Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.


If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

See Side Panel For Additional Precautions

NET CONTENTS:

AgrEvo Environmental Health

95 Chestnut Ridge Road Montvale, NJ 07645



Same Mac.

PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Avoid contamination of feed and foodstuffs. In the home, all food processing surfaces should be covered during treatment or thoroughly washed before use. Remove pets and cover or remove fish aquariums before spraying.

Environmental Hazards

This pesticide is extremely toxic to fish. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

80 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON EDIBLE CROPS.

SHAKE WELL BEFORE EACH USE.

Remove protective cap. Aim spray opening away from people. Push button to spray.

INDOORS

For House Flies, Mosquitoes and Gnats: Close all doors and windows. Point nozzle upward. Direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the goom with mist, then leave the freated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

> INSIDE - FOR CRAWLING INSECTS: Roaches, Acillarwigs, Grain Mites, Sowbugs, Spiders and Centipedes: Spray thoroughly into hiding places sheh as cracks, crevices, moist areas, openings

70

. .:

20/2

around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible.

OUTDOORS

OUTDOORS - TO KILL FLYING INSECTS:
Spray areas infested with Mosquitoes, House Flies,
Gnats (Midges), Biting Flies and Flying Moths
(Millers): Follow precautions listed above. Hold
container 18 to 24 inches from shrubs, bushes and
grassy areas. Spray these areas with a slow sweeping
motion at the rate of from 1 to 3 seconds per cubic
yard of space while moving away from the treated
area. To reduce annoyance from these flying insects,
treat open porches and patios in a similar manner.
Spray only when the air is still. Allow spray to
dissipate before occupying the treated area. Repeat
as necessary.

OUTDOORS - TO PROTECT ORNAMENTAL PLANTS: To control Aphids, Exposed Thrips, Adult White Flies, Leafhoppers, Plant Bugs, Spittlebugs and Japanese Beetles on similar rhododendrons, pines, spruces, hemlocks, spireas, dogwoods, palms, mimosas, toyons, viburnums, euonymus, oaks, maples, locusts, forsythias and altheas and Mimosa Webworm Larvae on mimosas, spray directly on insects. Hold container 18 inches from the infested plants, spray in short spray bursts of 1 to 3 seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

Spray peonies, irises, zinnias, snapdragons, ageratums, chrysanthemums, nasturtiums, daisies, geraniums, marigolds, daphnes, fatshederas, ivy, tobiras, queen sagos and broadleaf acacias for control of Aphids, Exposed Thrips, Adult White Flies, Leafhoppers, Plant Bugs, Spittlebugs and Japanese Beetles. Spray as directed above.

Ornamental quinces, junipers, cypress, ornamental mulberries, ash, cotoneasters, roses, camellias, azaleas, pyrecarthas and caenothus may also be treated to control the above pests but use with extreme caution. Certain varieties have shown some leaf burn under certain conditions of use such as heavy dosage. Repeat as necessary to obtain control.

OUTDOORS - TO PROTECT HOUSE PLANTS: Spray infested African violets, philodendrons, geraniums, begonias, coleus for control of listed insects. Repeat application as necessary. OUTDOORS - TO CONTROL NUISANCE PLANTS:

Ants: Spray trails, nests and points of entry. spray on ants where possible. Repeat as necessary.

Wasps: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

Spiders: Direct short bursts of spray on spiders and repeat as necessary.

STORAGE & DISPOSAL

Store in a cool, dry area. Do not transport or store below 32°F. Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.