

PYRENONE SMALL ANIMAL & KENNEL INSECTICIDE

EMULSIFIABLE CONCENTRATE

EPA Reg. No. 4816-406

A PYRENONE Three trafe

For Control of Type I had I you and Far Mites on Pogs and Cats

Projects Against Chick to Freets Bring Flies, and Mosquitoes.

The insecticidal mass hours of the one contrate are highly and rapidly effective a arrowing of the Contrate is designed for dilution with water as specified rates for use on animals or in kennels as a specific or it as my spray.

ACTIVE INGREDIENTS:

Pyrethrins

100.00

*Equivalent to 8.47% of (buty1carbity1) (b-propy piperony1) ether and to 2.37% of related compounds

CAUTION Keep Out Of Reach Of Children

See rear panel for additional precautions

PYRENONE - Registered Trademark of FMC Corporation

NET CONTENTS: 1 QUART

Mondoclused by

INDUSTRIAL DEPARTMENT

Niagara Chemical Division Middleport, N.Y. 14105

DIRECTIONS

I'thation of this concentrate should be made shoully before use in the quantity needed for the individual application. Water dilutions should not be kept for future use.

To Control Insects on Animals

Dilute this concentrate at the rate of 7 fluid ounce, in 1 gallon with water or 1 fluid ounce in 18 fluid ounces of water.

For rapid control of ticks, fleas and lice on dogs, and cats apply this dilution as a mist spray. To insure thorough coverage, part the fur by stroking against the lay of the hair as spray is applied. Spray directly on ticks. Apply to moisten the fur lightly.

To control spinose ear ticks and car mites, syringe the ears with a few drops of this dilution so that it can penetrate the the wax down to the surrounding tissues. Repeat treatment once or twice at 2 to 3 days intervals as necessary.

For temporary protection from mosquitoes, eye gnats, biting flies, and chiggers apply to cover the surface of the hair. Repeat treatment as necessary when insects are troublesome.

To Control Lisects in Kennels and Animal Quarters

Dilute at the raw of 13 fluid ounces in 1 gallon with water or 2 fluid ounces in 18 fluid ounces of water.

Removements, birds, and cover fish aquariums before spraying.

For rapid cleanout of <u>fleas</u> and <u>ticks</u> in kennels apply as a surface spray in a good machine adjusted to deliver coarse droptests. Apply to the point of run-off with special attention to cracks and crevices and other insect hiding places. Bedding should be removed and sprayed thoroughly.

To control <u>flies</u>, <u>mosquitoes</u> and <u>gnats</u> in kennels, apply this dilution in a good sprayer adjusted to deliver a fine mist. Use at the rate of approximately I ounce for each 1000 cubic feet of space. Fill the enclosed area with mist. Wherever practical, keep doors and windows closed for 10 minutes or more after application. Do not remain in treated areas and ventilate after treatment is completed.

CAUTION: Harmful if swallowed. Avoid contamination of food, feed, and utensils. This product is toxic to fish. Keep out of lakes streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Dispose of containers by rinsing with solvent, washing thoroughly or steam cleaning and discarding, disposing of wash water with normal industrial wastes.

Avoid contamination of food and utensils by keeping them, well covered while spraying the area.

043

Printed in U.S.A.

