



VAPORIZER CONCENTRATE 0.5-0.1

EPA Reg. No. 4816-410

An Effective Semi-Concentrated Pyrenone δ Insecticide.

For application in Tufa, Challenger, Tomado, Dina-Fog, Mircosol, Halaby, Klip-On, Commander, Ees-Kill and similar conventional thermol, mechanical, or compressed air fogging equipment.

For use in Bakeries, Food Plants, Restaurants, Hotels, Motels, Hospitals, Schools, Office Buildings and Food Processing or Handling Plants.

ACTIVE INGREDIENTS

Pyrethrins	0.1%
*Piperonyl Butoxide, Technical	0.5%
**INERT INGREDIENTS:	99.4%
	100.0%

*Equivalent to 0.4% of (butylcarbityl) (6-propylpiperonyl) ether and 0.1% of related compounds.

**Contains Petroleum Distillate

PYRENONE is a registered trademark of FMC Corporation

CAUTION

Keep Out Of Reach Of Children

See rear panel for additional precautions

Net Contents 1 Gallon



Manufactured by
INDUSTRIAL DEPARTMENT
Niagara Chemical Division
Middleport, N. Y. 14105

DIRECTIONS

Close windows and doors and shut off ventilating systems. Using appropriate fogging or vaporizing applicators, apply at rate of 3 to 6 ounces per 1000 cubic feet of space. The lower rate may be used at machine settings which provide finer particles of spray mist. Manufacturer's recommendations may often be helpful. The dosage and distribution should be sufficient to thoroughly fill the enclosure with vapor or fog. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of flies, including fruit flies, mosquitoes, silverfish, flying moths, gnats, crickets, roaches, spiders, scorpions, wasps, hornets, clover mites, cheese mites, cheese skippers, and these common stored product pests: granary weevils, rice weevils, grain beetles, spider beetles, cigarette beetles, Drugstore beetles, flour beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, Tobacco moths, meal worms, and grain mites.

CAUTION: Harmful if swallowed. Avoid contamination of feed or foodstuffs.

The use of this product as a space spray in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Do not remain in treated areas and ventilate the area after treatment is completed.

This insecticide is an oil base material and should not be sprayed, stored or otherwise exposed to an open flame. Not more than one gallon should be sprayed in 40,000 cubic feet of enclosed space in order to avoid an explosion hazard.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place or return to drum reconditioner.

052

Printed in U. S. A.

This document is for informational purposes only and should not be used for Phase 2
of the project without a change in status.

