[4816-414] D

DIRECTIONS

Hold container upright with nozzle away from you. Press valve down and spray as directed. Keep nozzle at least 30 inches from surfaces or fabrics to avoid possible staining.

Remove pets, birds and cover fish aquariums before spraying.

Flies, Mosquitoes, Gnats, Wasps, and Flying Moths: Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

Roaches, Water Bugs, Silverfish, Crickets, Spiders, Centipedes, and Sow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests, and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moth, Angoumois Grain Moth: Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

Flour Beetles, Weevils, Grain Mites: Spray into cracks and crevices, processing, and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings, and floors. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pers are kept. Repeat as necessary. Use only a recommended per spray on pers.

CAUTION: HARMFUL IF SWALLOWED. Avoid contamination of feed and foodstuffs. Vacate premises after treatment and ventilate before reoccupying.

When using this product as a space spray in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

WARNING: Contents under pressure. Do not puncture. Do not expose container to temperatures above 120°F. Never throw container into fire or incinerator.



EPA Reg. No. 4816-414

Contains PYRENONE

Industrial Strength - Low Dosage Required

For use in Schools, Hospitals, Institutions, Dairies, Supermarkets, Hotels, Restaurants, Drive-Ins, Metels, Drug Stores, Food Plants, and Animal Quarters.

Procedes Rapil Convert of Plie : Mergantoes, Grats, Wasps, Phone, Mothe (Miller, on the essible spaces of Roaches, Ants, Water Buss, Sile on he liber Beetle : Worlds, Grain Mites, and Spiders.

ACTIVE INGREDIENTS:

| Pyrethrins | |
|--------------------------------|-------|
| *Piperonyl Butoxide, Technical | |
| Petroleum Distillate | 15.5% |
| INERT INGREDIENTS | 80.0% |

*Equivalent to 3.2% of (butylcarbity1) (6-propylpiperony1) ether and to 0.8% of related compounds.

CAUTION KEEP OUT OF REACH OF CHILDREN.

SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

PYRENONE is a registered trademark of FMC Corporation

NET CONTENTS: 1 POUND 4 OUNCES



Manchactered by INDUSTRIAL DEPARTMENT

Niagara Chemical Division Middleport, N Y 14105