CODE 775.00

# DAIRY & LIVESTOCK SPRAY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

Harmful If swallowed. Avoid contamination of feed and foodstuffs. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of spray equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Reseal container and offer for reconditioning or triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe biplace. Consult Federal, state or local disposal authorities for approved al-Miternative procedures.

12500 D 518

A Ready-To-Use PYRENONE® Insecticide

#### KILLS AND REPELS

Kills Houseflies, Stable Flies, Horse Flies, Horn Flies, Mosquitoes, Gnats

Acceptable for Use on Dairy and Meat Animals and in Dairy Rooms, Hilk Rooms and Dairies

#### ACTIVE INGREDIENTS:

Pyrethrins . . . . 0.10% \*Piperonyl Butoxide, Technical 1.00≵ Petroleum Distillate . . . . 98.90%

\*Equivalent to 0.8% (butylcarbityl) (6propylpiperonyl) ether and 0.2% related compounds

PYRENONE - Registered trademark of Fairfield American Corp.

## Keep Out Of Reach Of Children


## CAUTION

Harmful if swallowed. See side panel for additional precautionary statements.

EPA Rea. No. 4816-433 EPA Est. No. 279-NY-1

NET CONTENTS. \_GALLONS NET CONTENTS. LITERS NET CONTENTS POUNDS NET CONTENTS\_ KILOS

BATCH



horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the tips of the hair. To control stable flies, horse flies, deer flies, apply 2 ounces per adult animal, sufficient to wet the hair, but not to soak the hide. Repeat treatment once or twice daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply the spray lightly to infested areas, using a stiff brush to get the spray to the base of the hair. Use 2 ounces or less per animal. Repeat every two to three weeks as required.

BARNS, HILK ROOMS, DAIRY PLANTS: To control flies, mosquitoes, and gnats, apply as a foo or fine mist, directing the spray toward the ceiling and upper corners of the room until the area is filled with mist, using about 2 ounces per 1000 cubic feet of space. For best results, close windows and doors before spraying and keep closed for ten to fifteen minutes. Vacate the area and ventilate before reoccupying. Repeat application as necessary.









Buyer assu es all risks of use, storage or handling of this material mut in strict accordance with directions diver 4-6.11

FAIRFIELD **AMERICAN** CORPORATION 3932 Salt Boad Medina, New York 14103