

[4816-440]

DIRECTIONS

SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

Flies, Mosquitoes, Gnats, Wasps, and Flying Moths: Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

Roaches, Water Bugs, Silverfish, Crickets, Spiders, Centipedes, and Sow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains, and plumbing, hitting insects with spray when ever possible. Repeat treatment as necessary.

Ants: Spray trails, nests, and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moth, Angoumois Grain Moth: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets, and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

Confused Flour Beetles, Saw-Toothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings, and floors. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Avoid contamination of feed and foodstuffs. When using this product in food processing plants, food should be removed or covered during treatment. All food processing surface should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

WARNING! Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.

INDUSTRIAL AQUEOUS PRESSURIZED 2.0-0.4

EPA Reg. No. 4816-440

A PYRENONE® Insecticide

Industrial Strength- Low Dosage Required

For Use in Schools, Hospitals, Institutions, Dairies, Hotels, Supermarkets, Restaurants, Drive-Ins, Motels, Drug Stores, Food Plants and Animal Quarters.

Provides Rapid Control of Flies, Mosquitoes, Gnats, Wasps, Flying Moths (Millers) and accessible stages of Roaches, Ants, Water Bugs, Silverfish, Flour Beetles, Weevils, Grain Mites, and Spiders.

ACTIVE INGREDIENTS

Pyrethrins 0.40%

*Piperonyl Butoxide, Technical 2.00%

Petroleum Distillate 7.40%

INERT INGREDIENTS 90.20%

*Equivalent to 1.6% of (butylcarbityl) (6-propylpiperonyl) ether and 0.4% of related compounds.

PYRENONE- FMC trademark

CAUTION

Keep Out Of Reach Of Children

See rear panel for additional precautions

CAUTION: Avoid breathing of vapor or contact with eyes or skin. Vacate premises after treatment and ventilate before reoccupying.

Remove pets, birds and cover fish aquariums before spraying.

NET WEIGHT: 1 POUND 4 OUNCES

MANUFACTURED IN U.S.A. BY

FMC CORPORATION

A.C.D. INDUSTRIAL SALES DEPARTMENT

FMC