DIRECTIONS

SHAKE BEFORE USING AND AT INTERVALS DURING PRO-LONGED USE.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

Flies, Mosquitoes, Gnats, Wasps, and Flying Moths: Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary

Roaches, Water Bugs, Silverfish, Crickets, Spiders, Centipedes, and Sow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains, and plumbing, hitting insects with spray when ever possible. Repeat treatment as necessary.

Ants: Spray trails, nests, and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moth, Angoumois Grain Moth: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets, and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

Confused Flour Beetles, Saw-Toothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites: Spray into cracks and crevices, processing and handling equipment, aroundcartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tults and seams. Take beds apart and spray into all joints. Treat base boards, moldings, and floors. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

CAUTION: Avoid breathing of vapor or contact with eves or skin. Vacate premises after treatment and ventilate before reoccupying.

Avoid contamination of feed and foodstuffs. When using this product in food processing plants, food should be removed or covered during treatment. All food processing surface should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

WARNING: Contents under pressure. Do not puncture. Do not expose container to temperature above 120°F. Never throw container into fire or incinerator.



INDUSTRIAL AQUEOUS PRESSURIZED 4.0—0.5

EPA Reg. No. 4816-441

A PYREONE "Insecticate

Industrial Strength Low Dosage Required

For Use in Schools, Hospitals, Institutions, Dairies, Hotels, Supermarkets, Restaurants, Drive-Ins, Motels, Drug Stores, Food Plants and Animal Quarters.

Provides Rapid Control of Flies, Mosquitoes, Gnats, Wasps, Flying Moths (Millers) and accessible stages of Roaches, Ants, Water Bugs, Silverfish, Flour Beetles, Weevils, Grain Mites, and Spiders?

*Equivalent to 3.2% of (butylcarbity1) (6-propylpiperony1) ether and 0.8% of related compounds.

CAUTIONKeep Out Of Reach Of Children

See rear panel for additional precautions

PYRENONE is a registered trademark of FMC Corporation

