

After washing and cleaning up the inside of the processing plant and just prior to bringing fruit into it, the entire space inside the cannery should be sprayed at a dilution of 6 parts concentrate to 50 parts water (6 pints with 7 gallons 3 pints water or 12 tablespoonsful with 3 1/2 pints water). At the rate of 1 gallon to 750 square feet (1 quart to 188 square feet) direct spray on walls, ceilings, and floors, paying special attention to forcing spray into all cracks and crevices.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Contact a physician or Poison Control Center. Do not induce vomiting unless directed by a physician. Vomiting may cause aspiration pneumonia.

If Inhaled: Remove affected person to fresh air. Apply artificial respiration if indicated.

If In Eyes: Flush eyes with plenty of water. Contact a physician if irritation persists.

If On Skin: Remove contaminated clothing and wash contaminated areas of skin with soap and water. Contact a physician if irritation persists.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Remove pets, birds and cover fish aquaria before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not apply as a space spray while food processing is underway. Except in Federally inspected meat and poultry plants, when applied as a surface spray with care and in accordance with the directions and precautions given above, food processing operations may continue. Do not apply at any time as an undiluted space spray in edible product areas of Federally inspected meat and poultry plants. Foods should be removed or covered before treatments. Thoroughly wash all food processing surfaces before reuse. In food processing plants and Federally inspected meat and poultry plants, all surfaces in processing areas must be washed and rinsed in potable water after spraying.

When using in animal quarters, do not apply directly to food, water or food supplements. Wash teats of dairy animals before milking.


ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour or spill near heat or open flame.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



BEST AVAILABLE COPY