

[4816-455]



FELINE INSECT SPRAY

EPA Reg. No. 4816-455

DIRECTIONS

To control fleas and lice on cats: Holding can upright, spray cat with short bursts from a distance of 10 to 12 inches. Starting at the tail, stroke against the lay of the hair, spraying the parted fur directly behind the hand to insure penetration of the coat. Dampen the fur up to the head. Spraying 20 to 30 seconds is usually sufficient for an adult cat. Repeat application when reinfestation occurs.

Do not treat or cause exposure of kittens under four weeks of age.

For best results in controlling tick infestations, locate ticks and spray directly on them.

To control infestations in sleeping quarters, after treating animals, spray bedding, walls and flooring with particular attention to cracks and crevices.

CAUTION: Avoid inhalation of mist. Avoid contamination of feed, foodstuffs or utensils. Wash hands after use.

Warning: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container in fire or incinerator.

For Control of Fleas, Lice and Ticks on Cats

A Non-Aqueous PYRENONE[®] Insecticide

ACTIVE INGREDIENTS:

Pyrethrins06%
*Piperonyl Butoxide, Technical60%
Petroleum Distillate	4.34%
INERT INGREDIENTS:	95.00%

*Equivalent to 0.48% (butylcarbityl) (6-propylpiperonyl) ether and to 0.12% of related compounds

CAUTION

Keep Out Of Reach Of Children

See rear panel for additional precautions

Net Weight _____

Pyrenone -Registered Trademark FMC Corporation

Manufactured by
INDUSTRIAL DEPARTMENT

Niagara Chemical Division
Middleport, N. Y. 14106



ACCEPTED

JAN 1 1975

U.S. DEPARTMENT OF AGRICULTURE
NATIONAL ANIMAL HEALTH LABORATORY
MIDWEST DIVISION
MIDDLEPORT, N.Y.

4816-455

SPECIMEN LABEL

Printed in U.S.A.