

CODE 724.30

PYRENONE® GARDEN SPRAY CONCENTRATE

A PYRENONE Insecticide Concentrate

Dilute with Water for Use on Vegetables and Ornamental Plants

Quick-Acting - Leaves No Harmful Residues

Acceptable for Use To Day of Harvest on Edible Plants

ACTIVE INGREDIENTS

Pyrethrins	1.00
*Piperonyl Butoxide, Technical	10.00
Petroleum Distillate	79.00
INERT INGREDIENTS	10.00
	100.00

*Equivalent to 5.0% (butylcarbityl)(6-propyl-piperonyl) ether and 2.0% related compounds

PYRENONE - Trademark of Fairfield American Corporation

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Warning: Harmful if swallowed

See side panel for additional precautionary labeling.

EPA Reg. No. 4816-462
EPA Est. No. 277-NY-1

BATCH

ACCEPTED

FEB 27 1978

When the Federal Insecticide, Fungicide, and Rodenticide Act is amended to the present date, the date of the amendment is 4816-462.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

This insecticide may be used outdoors or in greenhouses throughout the entire growing season or as a preharvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named on the plants listed below, apply concentrate at a rate of 12 to 36 fluid ounces per acre, in sufficient water for thorough coverage (normally 100 gallons per acre). For smaller gardens, use 0.6 to 1.5 fluid ounces of concentrate in 5 gallons of water or 1 to 3 tablespoons per 1 gallon. Apply to insure thorough coverage of upper and lower leaf surfaces.

Prepare only the amount of spray required. Hold-dilute spray more than 24 hours is not recommended.

CROP	TO CONTROL
ASPARAGUS	Asparagus Beetle
BEANS	Aphids, Leafhoppers, 12-Spotted Cucumber Beetle, White Fly, Mexican Bean Beetle
BROCCOLI	Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback
CABBAGE	Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug
CAULIFLOWER	Green Peach Aphid, Leaf Tier, Cabbage Looter
CELERY	Fireworms, Leafhopper
CRANBERRIES	Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid
EGGPLANTS	Cabbage Looter, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm
LETTUCE	Cabbage Looter, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm
MUSTARD GREENS, KALE, TURNIPS, COLLARDS	Cabbage Looter, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm
PEPPERS	Green Peach Aphid
POTATOES	Aphids, Colorado Potato Beetle, Flea Beetle, Leafhoppers
RADISH	Aphids, Flea Beetle
SQUASH	Aphids, Cabbage Looter, Cabbageworm
TOMATOES	Green Peach Aphid, Stink Bug, Colorado Potato Beetle, Flea Beetle
ORNAMENTALS	Africa Mites, Aster, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwood, Geraniums, Gladioli, Marigolds, Petunias, Rubber Plants, Wandering Jew, etc. Aphids, Flea Beetles, Leafhopper

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



FAIRFIELD
AMERICAN
CORPORATION

3032 Salt Road
Medina, New York 14103

2062

CODE 724.30

PYRENONE® GARDEN SPRAY CONCENTRATE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Reseal container and offer for reconditioning or triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place. Consult Federal, state or local disposal authorities for approved alternative procedures.

This insecticide may be used outdoors or in greenhouses throughout the entire growing season or as a preharvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.
To control insects named on the plants listed below, apply concentrate at a rate of 1/4 to 1/2 fluid ounces per acre, in sufficient water for thorough coverage (normally 100 gallons per acre). For smaller gardens, use 0.6 to 1.5 fluid ounces of concentrate in 5 to 15 gallons of water for 1 to 3 applications per 1 gallon. Apply to insure thorough coverage of upper and lower leaf surfaces.

A PYRENONE Insecticide Concentrate

Dilute With Water for Use on Vegetable and Ornamental Plants

Quick-Acting - Leaves No Harmful Residues

Acceptable for Use To Day of Harvest on Edible Plants

ACTIVE INGREDIENTS:

Pyrethrins	1.0%
Piperonyl Butoxide, Technical	10.0%
Petroleum Distillate	79.0%
INERT INGREDIENTS:	10.0%
	100.0%

Equivalent to 8.0% (butylcarbityl) (6-propylpiperonyl) ether and to 2.0% related compounds

PYRENONE - Trademark of Fairfield American Corporation

Keep Out Of Reach Of Children

CAUTION

Harmful if swallowed.
See side panel for additional precautionary statements.

EPA Reg. No. 4816-462
EPA Est. No. 279-NY-1

NET CONTENTS _____ GALLONS
NET CONTENTS _____ LITERS
NET CONTENTS _____ POUNDS
NET CONTENTS _____ KILOS


BATCH

Prepare only the amount of spray required. Holding dilute spray more than 24 hours is not recommended.

CROP	TO CONTROL
ASPARAGUS	Asparagus Beetle
BEANS	Aphids, Leafhoppers, 12-Spotted Cucumber Beetle, White Fly, Mexican Bean Beetle
BROCCOLI	Aphids, Cabbage Looper, Cross-striped Cabbage-worm, Diamondback Moth
CABBAGE	Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug
BRUSSELS SPROUTS	
CAULIFLOWER	
CELERY	Green Peach Aphid, Leaf Tier, Cabbage Looper
CRANBERRIES	Fireworms, Leafhopper
EGGPLANTS	Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid
LETTUCE	Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm
MUSTARD GREENS	Imported Cabbageworm, Diamondback Caterpillar, Aphids, Cabbage Looper
KALE	
COLLARDS	
TURNIPS	
PEPPERS	Green Peach Aphid
POTATOES	Aphids, Colorado Potato Beetle, Flea Beetle, Leafhoppers
RADISH	Aphids, Flea Beetle
SPINACH	Aphids, Cabbage Looper, Webworm
TOMATOES	Green Peach Aphid, Stink Bug, Colorado Potato Beetle, Flea Beetle
ORNAMENTALS	(African Violets, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwood, Geraniums, Gladioli, Marigolds, Roses, Rubber Plants, Wandering Jews, etc.) Aphids, Flea Beetle, Leafhopper

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

ACCEPTED
FEB 7 1978



FAIRFIELD AMERICAN CORPORATION

3932 Salt Road
Medina, New York 14103