

726.25

TETRALATE MULTI-PURPOSE INSECTICIDE E.C

Combining Two Highly Active Synthetic Pyrethroid Compounds

A Water Emulsifiable Insecticide Concentrate

For Use in Bakeries, Bottling Plants, Canneries, Flour Mills, Food Processing Plants, Hospitals, Hotels, Motels, Restaurants, Schools and on Ornamental Plants in Homes, Gardens and Greenhouses.

ACTIVE INGREDIENTS:

Tetramethrin	2.50%
Related compounds	0.34%
Resmethrin: [5-(phenylmethyl)-3-furanyl]-ethyl 2,2-dimethyl 3-(2-methyl-1-propenyl) cyclopropanecarboxylate	2.50%
Related compounds	0.46%
Aromatic petroleum hydrocarbons	20.00%
Petroleum Distillate	10.00%
INERT INGREDIENTS:	64.20%
	100.00%

1-cyclohexene-1,2-dicarboximidomethyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate

Cis/trans isomer ratio: Max. 30% (+) cis and min. 70% (+) trans.

Keep Out Of Reach Of Children

CAUTION

See rear panel for further precautions.

EPA Reg. No. 4816-470
EPA Est. No. 279-NY-1

NET CONTENTS _____ GALLONS
NET CONTENTS _____ LITERS
NET CONTENTS _____ KILOS
NET CONTENTS _____ POUNDS

BATCH _____

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Wash all contaminated clothing with soap and hot water before reuse. Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

Do not apply outdoors when weather conditions favor drift from the area. This product is toxic to fish and birds. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

FLAMMABLE! Keep away from heat and open flame.

Water dilutions should be prepared at room temperature. Exposure to low temperatures may cause cloudiness or crystallization. Prepare only the amount required for immediate use.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Reseal container and offer for reconditioning or triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place. Consult Federal, State or local disposal authorities for approved disposal procedures.

SEE REVERSE SIDE



**FAIRFIELD
AMERICAN
CORPORATION**

3932 Salt Road
Medina, New York 14103

19

BEST DOCUMENT AVAILABLE

The active components of this insecticide concentrate are the new synthetic pyrethroid compounds, tetramethrin and resmethrin. They provide quick knockdown and broad spectrum control.

This concentrate should be diluted with water before use, at a rate of 1 part concentrate to 9 parts water. The diluted spray should be applied in conventional mechanical or compressed air spraying or fogging equipment. It is not recommended for use in thermal generating equipment.


To control roaches, ants, spiders, silverfish, crickets, centipedes, earwigs, sowbugs and grain mites: Apply as a coarse droplet spray directed into cracks, crevices and other hiding places along baseboards, around the base of machinery and under pallets. Treat all returned bottle cartons and cases. Spray thoroughly contacting insects directly whenever possible. Spray badly infested areas daily. In other areas repeat treatments twice weekly or when reinfestation occurs.

To control flies, mosquitoes, gnats and small flying moths: Close doors and windows and shut off ventilating systems. Apply as a fog or fine mist directing the spray toward the upper portions of the room and towards windows and other light sources which attract these insects. Thoroughly fill the area with mist using $\frac{1}{2}$ to 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Vacate after treatment and ventilate before reoccupying. Repeat treatment as needed.

ON ORNAMENTAL PLANTS: To control aphids, two-spotted spider mites and adult white flies on ornamental house and garden plants (African violet, azalea, Christmas cherry, chrysanthemum, cineraria, dieffenbachia, fuschia, gloxinia, ivy, lantana, Neanth bella palm, petunia, rose and snapdragon and on outdoor plants such as azalea, chrysanthemum, euonymus, ivy, marigold, petunia, rose, snapdragon and zinnia). Apply as a fog or fine mist, spraying plants with a sweeping motion so that the mist drifts into the plants. Direct spray so that the upper and lower portions of leaves are contacted. Use caution on new growth and tender foliage. When using outdoors, spray when air is still. Repeat as necessary.

To control tent caterpillars, canker worms, garden web worms, and the above pests on ornamental crabapple, cherries and plum trees spray as above. Do not spray bearing fruit trees.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



10-30-80

000000

BEST DOCUMENT AVAILABLE