



# PYRENONE<sup>®</sup> 1-0.2 FOOD PLANT SPRAY

EPA Reg. No. 4816-471

A Ready-To-Use PYRENONE<sup>®</sup> Insecticide

Designed For Use in Edible Product Areas of Meat and Poultry Processing Plants

The insecticidal components piperonyl butoxide and pyrethrins are classified as food additives, and when this product is used as directed allowable tolerance levels will not be exceeded.

## ACTIVE INGREDIENTS:

Pyrethrins	0.20%	100%
*Piperonyl Butoxide, Technical	1.00%	
Petroleum Distillate	98.80%	

\*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% of related compounds

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN.**

See rear panel for additional precautions

**Net Contents 1 Gallon**

PYRENONE is a registered trademark of FMC Corporation



Manufactured by  
INDUSTRIAL DEPARTMENT  
**Niagara Chemical Division**  
Middleport, N. Y. 14105

## DIRECTIONS

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

**SPACE SPRAY:** For rapid control of houseflies, fruit flies, mosquitoes, flying moths and gnats: Close doors and windows and shut off ventilating systems. Apply using a good sprayer adjusted to deliver a fine mist. Direct spray toward the upper portions of the enclosure, filling the room with mist. Use ½ to 1 ounce per 1000 cubic feet of space.


To kill exposed stages of roaches, ants, spiders, clover mites, cheese mites, confused flour beetles, saw-toothed grain beetles, red flour beetles, rice weevils, granary weevils, and grain mites: Treat as above using 2 to 3 ounces per 1000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

**SURFACE SPRAY:** To insure maximum control of the crawling insects named above, apply as a coarse mist directed toward insect, into cracks and crevices, along baseboards, around the base of machinery, containers of stored products and other areas where insects may hide. Spray ant trails, nests and points of entry. Spray badly infested areas daily. In other areas, repeat treatment twice weekly or when reinfestation occurs.

**CAUTION:** Harmful if swallowed. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label.

Do not reuse can containing this insecticide. Destroy by perforating or crushing and burying in a safe place. Drums may be reused after thorough cleaning.



4816-471

03/12/1973

1/1