# AQUEOUS Pyrenone® GARDEN SPRAY

An effective Insecticide for Use on Vegetable and Ornamental Pests

A Concentrated Pyrenone Micro-Emulsion, Clear Stable

Apply with Conventional Mechanical Compressed Air Equipment

ACTIVE INGREDIENTS:
Pyrethrins............................. 0.02%
*Piperonyl Butoxide, Technical........ 0.20%
Petroleum Distillate.................. 0.08%
INERT INGREDIENTS: .................... 99.70%
100.00%

*Equivalent to 0.16% (butylcarbityl)(6-propyl-piperonyl) ether and 0.04% related compound.

## Keep Out Of Reach Of Children

## CAUTION

See side panel for additional precautions.

PYRENONE – Registered trademark of Fairfield American Corporation

EPA Reg. No. 4816-482
EPA Est. No. 279-NY-1

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

NET CONTENTS_____

| CROP | INSECTS CONTROLLED |
|------|--------------------|
| Asparagus | Asparagus Beetles |
| Beans | Aphids, Leafhoppers, White-flies, Mexican Bean Beetles, 12-Spotted Cucumber Beetles |
| Broccoli, Cauliflower, Cabbage, Brussels Sprouts | Aphids, Cabbage Looper, Cross-striped Cabbageworm Diamond-back Moth larvae, Flea Beetle, Harlequin Bug, Imported Cab-bageworm, Stink Bug |
| Celery | Green Peach Aphid, Cabbage Looper, Leaftier |
| Cranberries | Fireworms, Leafhoppers |
| Eggplant | Blister Beetle, Colorado Potato Beetle, Flea Beetles, Green Peach Aphids |
| Lettuce | Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm |
| Mustard Greens, Kale, Collards, Turnips | Imported Cabbageworms, Dia-mondback Caterpiller, Aphids, Cabbage Looper |
| Peppers | Green Peach Aphid |
| Potatoes | Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers |
| Radishes | Aphids, Flea Beetle |
| Spinach | Aphids, Cabbage Looper, Webworm |
| Tomatoes | Green Peach Aphid, Colorado Potato Beetles, Stink Bugs, Whiteflies  Flea Beetles |
| African Violets | Aphids, Flea Beetles, Leafhoppers |
| Asters | Aphids, Flea Beetles, Leafhoppers, Whiteflies |
| Azaleas | Aphids, Flea Beetles, Leafhoppers |
| Begonias | Aphids, Flea Beetles, Leafhoppers |
| Camellias | Aphids, Flea Beetles, Leafhoppers |
| Carnations | Aphids, Flea Beetles, Leafhoppers, Whiteflies |
| Chrysanthemums | Aphids, Flea Beetles, Leafhoppers, Whiteflies |
| Dahlias, Dogwood | Aphids, Flea Beetles, Leafhoppers |
| Geraniums, Gladioli | Aphids, Flea Beetles, Leafhoppers, Whiteflies |
| Marigolds | Aphids, Flea Beetles, Leafhoppers, Whiteflies |
| Roses | Aphids, Flea Beetles, Leafhoppers |
| Wandering Jew | Aphids, Flea Beetles, Leafhoppers |

INDOORS AS A SPACE SPRAY
To control flies, fruit flies, mosquitoes, gnats and small flying moths: apply using a food space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treat-ment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4ounces per 1000 square feet).

To control mosquito larvae, apply at the rate of 775 gallons per acre or 18 gallons per 1000 square feet. Apply as a coarse wetting spray to the border or on shallow waters of wetlands, swamps and marshes where mosquito larvae is known to breed.

## FAIRFIELD AMERICAN CORPORATION
3932 Salt Road
Medina, New York 14103