

AQUEOUS Pyrenone® GARDEN SPRAY

CODE 724.00

Non-aerosol Pump

An effective Insecticide for Use on Vegetable and Ornamental Pests

A Concentrated Pyrenone Micro-Emulsion, Clear Stable

Apply with Conventional Mechanical Compressed Air Equipment

ACTIVE INGREDIENTS:

Pyrethrins.....	0.02%
*Piperonyl Butoxide, Technical.....	0.20%
Petroleum Distillate.....	0.08%
INERT INGREDIENTS:	99.70%
	100.00%

*Equivalent to 0.16% (butylcarbityl)(6-propylpiperonyl) ether and 0.04% related compound.

Keep Out Of Reach Of Children

CAUTION

See side panel for additional precautions.

DO NOT TRANSPORT OR STORE UNDER 32°F.

If freezing occurs, thaw and agitate vigorously

PYRENONE - Registered trademark of Fairfield American Corporation

EPA Reg. No. 4816-482

EPA Est. No. 279-NY-1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO OPERATE: Shake well before use. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

This insecticide may be used indoors, outdoors or in greenhouse throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

CROP	INSECTS CONTROLLED
Asparagus	Asparagus Beetles
Beans	Aphids, Leafhoppers, Whiteflies, Mexican Bean Beetles, 12-Spotted Cucumber Beetles
Broccoli, Cauliflower, Cabbage, Brussels Sprouts	Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback Moth larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug
Celery	Green Peach Aphid, Cabbage Looper, Leafminer
Cranberries	Fireworms, Leafhoppers
Eggplant	Blister Beetle, Colorado Potato Beetle, Flea Beetles, Green Peach Aphids
Lettuce	Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm
Mustard Greens, Kale, Collards, Turnips	Imported Cabbageworms, Diamondback Caterpillar, Aphids, Cabbage Looper
Peppers	Green Peach Aphid
Potatoes	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers
Radishes	Aphids, Flea Beetles
Spinach	Aphids, Cabbage Looper, Webworm
Tomatoes	Green Peach Aphid, Colorado Potato Beetles, Stink Bugs, Whiteflies, Flea Beetles

African Violets, Aphids, Flea Beetles, Azaleas, Begonias, Leafhoppers, Camellias, Dahlias, Dogwood, Roses, Wandering Jew

Asters, Carnations, Aphids, Flea Beetles, Chrysanthemums, Leafhoppers, Whiteflies, Geraniums, Gladioli, Marigold

INDOORS AS A SPACE SPRAY

To control flies, fruit flies, mosquitoes, gnats and small flying moths: apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and turn off ventilating systems. Direct spray toward portions of the enclosure filling

the room with mist. Use 4 to 5 ounces per 1000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4ounces per 1000 square feet).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or .18 gallons per 1000 square feet. Apply as a coarse wetting spray to the border or on shallow waters of wetlands, swamps and marshes where mosquito larvae is known to breed.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

CAUTION: Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.



ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.



**FAIRFIELD
AMERICAN**
CORPORATION
3012 SALT ROAD
MEDINA, N.Y.

4816-482
05/13/1982
CODE 724.00