

pg 102
Pm 16
EPA# 4816-483
CHF 1/30/90

DOSE SCALE

TOMATO & VEGETABLE INSECT SPRAY

EPA Reg. No. 4816-483
EPA Est. No. 279-NY-1

PROVIDES RAPID CONTROL OF INSECTS NAMED ON THE LABEL.
MAY BE USED ON FOOD CROPS TO WITHIN ONE DAY OF HARVEST.
CONTAINS TWO NATURAL INSECTICIDES:
PYRETHRINS AND ROTENONE (HOUSE' RESINS)

ACTIVE INGREDIENTS:

Pyrethrins	0.030%
*Piperonyl Butoxide, Technical	0.100%
Rotenone	0.100%
Other Associated Resins	0.100%
INERT INGREDIENTS	99.744%
	100.000%

*Equivalent to min. 0.100% (butylcarbitol) (n-propylpiperonyl) ether and 0.200% related compounds

Keep Out of Reach of Children

CAUTION

See side panel for additional precautions

NET CONTENTS _____

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label and apply as directed.

Shake well before using.

For best results, use throughout the growing season to prevent buildup of insect populations. Spray plants from a distance of 1 1/2 to 2 feet using short bursts (one second). To prevent injury to plants, avoid visibly wetting the foliage.

Apply in the early morning or late evening when the air is still, treating all foliage with particular attention to the undersides of the leaves. Contact insects directly whenever possible. Repeat applications at 7 to 10 day intervals or after heavy rains. Do not apply to food crops within one day of harvest.

USE ON: asparagus, beans, broccoli, cabbage, bean sprouts, cauliflower, celery, cranberries, grapes, cucumbers, melons, squash, eggplants, lettuce, mustard greens, kale, collards, turnips, beets, cots, peas, beans, radishes, spinach, strawberries, sweet corn, tomatoes and watermelon.
TO CONTROL: asparagus beetles, aphids, leafhoppers, whiteflies, Mexican bean beetles, 11-spotted cucumber beetles, cabbage loopers, corn-restricted dipterocorms, diamondback moth caterpillars, flea beetles, hornworms.

Not Reviewed. Registrant claims to be in accordance with
Registration of Pesticide Product Guidance for Phase 2
Response, Page 2.12, Active to Inert Change in Status.

BEST AVAILABLE COPY

Imported cabbageworms, stink bugs, green peach aphids, leafminers, fireworms, blister beetles, Colorado potato beetles, webworms, Japanese beetles, leaf beetles, leaf rollers, striped cucumber beetles, hornworms.
 USE ON ORNAMENTALS: African violets, lilies, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, dogwood, geraniums, gladioli, marigolds, roses, rubber plants, and wandering Jew.
 TO CONTROL: Aphids, flea beetles, leafhoppers, whiteflies, covey moth caterpillars and adults, tent and forest caterpillars, and adult Japanese beetles.

STORAGE AND DISPOSAL

Store in a cool, dry area. Do not transport or store below 32° F.

Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Keep out of reach of children. Harmful if swallowed or inhaled. Avoid breathing spray or contact with eyes or skin. Wash thoroughly after use. Do not contaminate feed and feedstuffs.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent areas. Do not contaminate water when disposing of equipment washwaters.


PHYSICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate containers. Exposure to temperatures above 130°F may cause bursting.

STATEMENT OF FIRST AID TREATMENT

IF ON SKIN: Wash with soap and water.
 IF IN EYES: Flush with water for 15 minutes. If irritation persists, consult a physician.
 IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

User assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein.



BEST AVAILABLE COPY