CODE 721.40

In EPA Leiter Patrick

TUTETONE 1-0.2 FOOD PLANT SPRAY

EPA Reg. No. 4816-471 EPA Est. No. 279-NY-1

A Ready-To-Use PYNERONE® Insecticide

Designed for Use in Edible Product Areas Of Meat And Poultry Processing Plants.

Food Additive Tolerances Have Been Istablished for The Insecticide Components In This Product. When This Product is Used As Directed, Allowable Tolerance Levels Will Not the Exceeded.

ACTIVE INGREDIENTS

| Pyrethrins | 0.20% |
|--------------------------------|-----------|
| *Piperonyl Butoxide, Technical | 1.00% |
| "INERT INGREDIENTS | 98.80% |
| | 100 00% |

*tourivalent to min. O B% (butylcarbity)) (6-propylpiperonyl) ether and 0.25 related compounds

"Contains Petroleum Distillate

Registered trademark of PYRELOGIC Fairfield American Curporation

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautions,

NET CONTENTS:

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SPACI. SPRAY

for rapid control of bouse files, fruit flies, mosquitoes, flying moths and gnats: Close doors and windows and shut off ventilating systems. Apply using a good sprayer adjusted to deliver a fine mist. Direct spray toward upper portions of the enclosure filling the room with mist, Use 1/2 to 1 ounce per 1000 cubic feet of space.


To kill exposed stages of cockroaches, ants, spiders, clover mites, cheese mites, confosed floor beetles, saxtoothed grain beetles, red floor beetles, rice weevils. granary weevils and grain mites. Treal as above using 2 to 3 ounces per 1000 cubic feet of space. Keep area closed for at least 15 minutes. Vucate treated area and ventilate before reoccupying, Repeat treatment when reinfestation occurs.

SURFACE SPRAY

To insure maximum control of the crawling insects named above, apply as a coarse mist freeted toward insects, into cracks and crevices, along baseboards, around the base of machinery, containers of stored products and other areas where insects may hide, Sprmy ant trails, nests and points of entry. Sprny badly infested areas daily. In other areas, repeat treatment twice weekly or when reinfestation occurs.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.



Harmful if swallowed. Avoid contamination of feed and foodstuffs. Do not apply as a space spray while food processing is underway. Except in Federally inspected meat and poultry plants, when applied as a surface spray with care and in accordance with the directions and precautions given mbove, food processing operations may continue. Foods should be removed or covered before treatments. All surfaces in processing areas must be washed and rinsed with potable water after spraying. Remove pets and birds and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

Il Swallowed: Contact a physician or Poison Control Center immediately, Do not Induce vomiting unless directed by a physician. Vomiting may cause aspiration pneumonia.

If Inhaled: Remove affected person to fresh alr. Apply artificial respiration if Indicated.


If In Eyes: Flush with plenty of water. Contact a physician if irritation persists,

11 on Skin: Remove contaminated clothing and wash affected areas of skin with soap and water.

EMVERORMENTAL HAZARDS

This product is toxic to fish be not apply directly to takes, streams or ponds. Donot contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.



BEST AVAILABLE COPY