

Apply at the rate of 30 pounds of dust per acre. Treat only when the insects are present and repeat as required to maintain control. Apply dust uniformly to upper and lower leaf surfaces.

Pyrenone Flexi-dust

CROP	TO CONTROL
ASPARAGUS	Asparagus Beetles
BEANS	Aphids, Leafhoppers, White Flies, Mexican Bean Beetles
BROCCOLI	Aphids, Cabbage Loopers, Diamond-back Moth Larvae, Cross-striped
CABBAGE	Cabbageworms, Flea Beetles, Harlequin Bugs, Imported Cabbageworms
BRUSSEL SPROUTS	
CAULIFLOWER	
CELERY	Green Peach Aphids, Leaf Tiers, Cabbage Loopers
CRANBERRIES	Fireworms, Leafhoppers
EGGPLANTS	Colorado Potato Beetles, Flea Beetles, Green Peach Aphids
LETTUCE	Cabbage Loopers, Green Peach Aphids, Diamondback Moth Larvae, Imported Cabbageworms
PEPPERS	Green Peach Aphids
POTATOES	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers
RADISH	Aphids, Flea Beetles
SPINACH	Aphids, Cabbage Loopers, Webworms
TOMATOES	Green Peach Aphids, Flea Beetles, Colorado Potato Beetles
ORNAMENTALS	Aphids, Flea Beetles, Leafhoppers on: African Violets, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwoods, Geraniums, Gladiolus, Marigolds, Roses, etc.

Fruit Fly Control of Fruits and Vegetables: To control fruit flies (*Drosophila*) on apples, blackberries, boysenberries, currants, dewberries, grapes, loganberries, peaches, pears, plums, melons and tomatoes, apply as a post-harvest application. Dust fruits and vegetables immediately after picking while in containers in the field. Remove crops from field as soon as they are picked and have been dusted. Do not hold picked crops in the field overnight. If the crops are held at the processing plants more than 20 hours, a repeat application should be made. Dust lightly but thoroughly.

Stored Grain Protectant: For use on corn, barley, rye, rice, buckwheat, birdseed and other seeds in storage.

Pyrenone Flexi-Dust kills accessible stages of many grain insects particularly rice weevils and granary weevils and protects against many insects in the bran bug group including flour beetles and grain beetles. One treatment is usually adequate for a full storage season.

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
*Piperonyl Butoxide, Technical	1.00%
Petroleum Distillate	1.90%
INERT INGREDIENTS:	97.00%
	100.00%

*Equivalent to 0.8% (butylcarbityl)(6-propylpiperonyl) ether and 0.2% related compounds

KEEP OUT OF REACH OF CHILDREN

CAUTION

Avoid Inhalation of dust.

See side panel for additional precautionary statements.

EPA Reg. No. 4816-496

EPA Est. No. 279-NY-1

PYRENONE - Registered trademark of Fairfield American Corporation

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid inhalation of dust.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of streams, lakes or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that can not be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

4816-496

2/9/1979

Page 132

292



Treated grains require only normal cleaning before processing for human food and may be fed to livestock without hazard. Shelled corn may require cleaning if subject to inspection for grading purposes.



Whole grains and seeds to be treated should be in condition for normal storage. Treatment should be applied when the grain goes into storage and before serious infestation has developed.

How Much Pyrenone Flexi-Dust to Apply:

Application on grains and seeds should be made at the rate of 80 to 100 pounds per 1,000 bushels (8/10 to 1 pound per 10 bushels). The dust should be sufficiently well mixed with the whole grain or seeds to assure even distribution. It can be applied to a moving stream of grain in the usual or typical conveyor. Small lots of grain or seeds can be treated by hand, with a shovel or other suitable equipment.

Pyrenone Flexi-Dust on ear corn including snapped corn should be applied at the same rate specified above and should be sufficiently well mixed with the corn to assure even distribution. One method is by dusting the protectant over alternate shallow layers of the corn.

Snapped (unshucked) corn should be shucked before treatment for best results. Otherwise the benefit of the treatment is primarily for the prevention of insect migration from infested ears. The amount of ear corn that will produce a bushel of shelled corn is figured as an "ear-corn bushel".



**FAIRFIELD
AMERICAN
CORPORATION**
3932 Salt Road
Medina, NY 14103

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

mc