

CODE 721.8

FOOD PLANT FOGGING INSECTICIDE

EPA Reg. No. 4816-512

EPA Est. No. 279-NY-1

A Concentrated PYRENONE® Product

DESIGNED FOR USE IN EDIBLE PRODUCT AREAS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER THE U.S. DEPARTMENT OF AGRICULTURE MEAT, POULTRY, RABBIT, SHELL EGG GRADING AND EGG PRODUCTS INSPECTION PROGRAMS

APPLY WITH EQUIPMENT DESIGNED AND ADJUSTED TO DELIVER A TRUE AEROSOL MIST.

ACTIVE INGREDIENTS:

Pyrethrins.....	0.5%
*Piperonyl Butoxide, Technical.....	5.0%
Petroleum Distillate.....	94.5%
	<u>100.0%</u>

*Equivalent to 4.0% (butylcarbityl)(6-propylpiperonyl) ether and 1.0% related compounds.

PYRENONE - Registered Trademark of Fairfield American Corporation

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See reverse side for additional precautions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

50.1.1

NET CONTENTS: 4.0.0.1

72180 12/85


The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment and thoroughly cleaned and rinsed with potable water, before using. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

SPACE SPRAY: For control of accessible, exposed stages of flies including fruit flies, mosquitoes, small flying moths, gnats, wasps, hornets, cheese skippers and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, meal worms and grain mites. Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. For best control of wasps and hornets, apply directly to nests in the evening when the insects have returned to the nests. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

SPACE AND CONTACT SPRAY: For control of accessible exposed stage of silverfish, crickets, cockroaches, spiders, scorpions, clover mites, cheese mites, earwigs and ants, first using equipment which can deliver a pin spray, direct spray into all cracks and crevice in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects are known to seek shelter. Contact as many insects as possible, spray directly all ant trails. Do allow spray to contact stored food. Then, closing all doors, windows and ventilating systems, proceed with space spraying direct as given above. Use a dosage rate of 1 ounce per 1000 cubic feet of space. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

IN MUSHROOM PRODUCTION AND PROCESSING: To control mushroom flies and fungus gnats in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulation fans should be left in operation, but for best results, refrigerators should be turned off. In packing and canning plants, use a rate of 1 ounce to 2000 cubic feet of space.

SEE REVERSE SIDE



STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, silt, earth, fuller's earth, etc. Dispose of with chemical waste.
PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.
CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.
CONTAINERS ONE GALLON AND SMALLER: Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment before application. Do not apply while food processing is under way. After space spraying in meat packing plants, bakeries and all other food processing plants wash all equipment thoroughly with a suitable detergent and rinse with potable water (benches, shelving, etc.), where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center immediately. Do not induce vomiting unless directed to do so by a physician. Aspiration is a potential hazard.

IF IN EYES: Flush with plenty of water. Contact a physician if irritation persists.

IF ON SKIN: Wash affected areas with plenty of soap and water.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

BATCH

Printed in U.S.A.

FOOD PLANT FOGGING INSECTICIDE

JAN 16 1986

Use in... Insecticide
Fungicide, and Herbicide And
as a fumigant for the pest
mentioned on the pesticide
registration label
EPA Reg. No. 4816-51

EPA Reg. No. 4816-512

EPA Est. No. 279-NY-1

A Concentrated PYRENONE® Product

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment and thoroughly cleaned and rinsed with potable water, before using. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

DESIGNED FOR USE IN EDIBLE PRODUCT AREAS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER THE U.S. DEPARTMENT OF AGRICULTURE MEAT, POULTRY, RABBIT, SHELL EGG GRADING AND EGG PRODUCTS INSPECTION PROGRAMS

SPACE SPRAY: For control of accessible, exposed stages of flies, including fruit flies, mosquitoes, small flying moths, gnats, wasps, hornets, cheese skippers and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, meal worms and grain mites. Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 70 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. For best control of wasps and hornets, apply directly to nests in the evening when the insects have returned to the nests. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

APPLY WITH EQUIPMENT DESIGNED AND ADJUSTED TO DELIVER A FINE AEROSOL MIST.

ACTIVE INGREDIENTS

Pyrethrins.....	0.5%
*Piperonyl Butoxide, Technical.....	5.0%
**INERT INGREDIENTS.....	94.5%
	100.0%

*Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and 1.0% related compounds.

**Contains Petroleum Distillate

PYRENONE - Registered trademark of Fairfield American Corporation.

SPACE AND CONTACT SPRAY: For control of accessible exposed stages of silverfish, crickets, cockroaches, spiders, scorpions, clover mites, cheese mites, earwigs and ants, first using equipment which can deliver a pin spray, direct spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects are known to seek shelter. Contact as many insects as possible, spray directly all ant trails. Do not allow spray to contact stored food. Then, closing all doors, windows and ventilating systems, proceed with space spraying directions as given above. Use a dosage rate of 1 ounce per 1000 cubic feet of space. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

IN MUSHROOM PRODUCTION AND PROCESSING: To control mushroom flies and fungus gnats in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigerators should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See reverse side for additional precautions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NET CONTENTS: 1.0 LITER

NET CONTENTS: 1.0 LITER

NET CONTENTS: 1.0 LITER

7/190 12/85

BEST AVAILABLE COPY

SEE REVERSE SIDE

**FAIRFIELD
AMERICAN
CORPORATION**
238 WILSON AVE
NEWARK, N.J. 07102