

EGGPLANTS	Colorado Potato Beetles, Flea Beetles, Green Peach Aphids
LETTUCE	Cabbage Loopers, Green Peach Aphid., Diamondback Moth Larvae, Imported Cabbageworms
PEPPERS	Green Peach Aphids
POTATOES	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers
RADISH	Aphids, Flea Beetles
SPINACH	Aphids, Cabbage loopers, Webworms
TOMATOES	Green Peach Aphids, Flea Beetles, Colorado Potato Beetles

Fruit Fly Control of Fruits and Vegetables:
To control fruit flies on apples, blackberries, boysenberries, currants, dewberries, grapes, loganberries, peaches, pears, plums, prunes, melons and tomatoes, apply as a post-harvest application. Dust fruits and vegetables immediately after picking while in containers in the field. Remove crops from field as soon as they are picked and have been dusted. Do not hold picked crops in the field overnight. If the crops are held at the processing plants more than 20 hours, a repeat application should be made. Dust lightly but thoroughly.

Stored Grain Protectant:

For use on corn, barley, rice, rye, buckwheat, birdseed, and other seeds in storage.

This product kills accessible stages of many grain insects, particularly rice weevils and granary weevils, and protects against many insects in the bran bug group, including flour beetles and grain beetles. One treatment is usually adequate for a full storage season.

Treated grains require only normal cleaning before processing for human food and may be fed to livestock without hazard. Shelled corn may require cleaning if subject to inspection for grading purposes.

Whole grains and seeds to be treated should be in condition for normal storage. Treatment should be applied when grain goes into storage and before serious infestation has developed.

How Much To Apply:

Application on grains and seeds should be made at the rate of 80 to 100 pounds per 1000 bushels (13 ounces to 1 pound per 10 bushels). The dust should be sufficiently well mixed with the whole grain seeds to assure even distribution. It can be applied to a moving stream of grain in the usual or typical conveyor. Small lots of grain or seeds can be treated by hand, with a shovel or other suitable equipment.

On ear corn this insecticide should be applied at the same rate specified above and should be sufficiently well mixed with the corn to assure even distribution. One method is by dusting the protectant over alternate shallow layers of the corn.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL:

Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL:

Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Avoid inhalation of dust.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

BATCH