

1/2 4816-518 RW 17

CODE 722-20

ACCEPTED
FEB 24 1988
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4816-518

AQUEOUS FOOD PLANT Pyrenone[®] FOGGING INSECTICIDE

EPA REG. NO. 4816-518

EPA EST. NO. 279-NY-1

A CONCENTRATED PYRENONE[®] MICRO EMULSION

- CLEAR
- STABLE
- LEAVES NO OILY OR OBJECTIONABLE RESIDUE

DESIGNED FOR USE IN EDIBLE PRODUCT AREAS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER THE MEAT, POULTRY, SHELL EGG GRADING AND EGG PRODUCTS INSPECTION PROGRAMS AND IN OTHER FOOD PROCESSING PLANTS.

APPLY WITH A CONVENTIONAL MECHANICAL OR COMPRESSED AIR EQUIPMENT DESIGNED AND ADJUSTED TO DELIVER A TRUE AEROSOL MIST.

ACTIVE INGREDIENTS:

PYRETHRINS.....	0.5%
*PIPERONYL BUTOXIDE, TECHNICAL.....	5.0%
INERT INGREDIENTS.....	94.5%
	100.0%

*Equivalent to 4.0% (butylcarbityl)(6-propylpiperonyl) ether and 1.0% related compounds.

PYRENONE - Registered Trademark of Fairfield American Corporation.

**KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS**

DO NOT TRANSPORT OR STORE UNDER 32°F

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment and thoroughly cleaned with an effective cleaning compound followed by a potable water rinse before using.

Close windows and doors and shut off ventilating systems. Use a non-thermal fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

IN BARN, POULTRY HOUSES AND STORAGE ROOMS. To control HOUSE FLIES, STABLE FLIES and HORN FLIES in barns, poultry houses, warehouses, storage bins and storage rooms: dilute 1 part of material with 4 parts of water. Apply as a fine mist over the livestock in stanchions or holding pens. Use a fogger or mist generator to direct spray above the animals and to the center of the area where animals are housed. Apply 1 ounce per 1000 cubic feet uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in dairy barns, ventilate the barn thoroughly before milking. Repeat daily in the morning and afternoon, or as required.

The above described application will kill accessible, exposed stages of FLIES, including FRUIT FLIES, MOSQUITOES, SILVERFISH, FLYING HORN, GNATS, CRICKETS, COCKROACHES, SPIDERS, SCORPIONS, WASPS, HERBETS, CLOSER MITES, CHEESE MITES, CHEESE SKIPPERS and these common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPINER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANOPLURIS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, MEAL MOTHS and GRAIN MITES.

IN MUSHROOM PRODUCTION AND PROCESSING. To control adult MUSHROOM FLIES, FLUGUS GNATS and CECID FLIES in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cubic feet of space.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

CONTAINERS (E.G. GALLON AND SWIGGERS): Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Do not pour or spill near heat or open flame. Remove pets, birds and cover fish aquaria before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water.

NET CONTENTS: _____

BATCH _____


SEE REVERSE SIDE

72220 1/86

Printed U.S.A.



**FAIRFIELD
AMERICAN**
CORPORATION
238 WILSON AVENUE
NEWARK, N.J. 07105



STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never give anything by mouth to an unconscious person.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

IF IN EYES: Flush eyes with plenty of water. Contact a physician if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.