

CODE 820.25

Reg # 4816-523 PM-17  
**Pyrenone<sup>®</sup> Dairy and Food Plant Aerosol**

Contains PYRENONE

Industrial Strength - Low Dosage Required

For Use in Non-Residential Food Handling Areas including Dairies and Milkrooms

KILLS House Flies, Mosquitoes and Gnats

**ACTIVE INGREDIENTS**

Pyrethrins.....	0.5%
*Piperonyl Butoxide, Technical.....	5.0%
**INERT INGREDIENTS.....	94.5%
	<u>100.0%</u>

\*Equivalent to 4 1/2 (butylcarbityl)(6-propylpiperonyl) ether and 1 1/2 related compounds

\*\*Contains Petroleum Distillate

PYRENONE - Registered trademark of Fairfield American Corporation

EPA Reg. No. 4816-523  
EPA Est. No. 279-NY-1

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

Harmful if swallowed. Do not apply over uncovered milk or food products.

See rear panel for additional precautions.

**ACCEPTED**

MAY 08 1979

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 4816-523 EPA Reg. No.

NET WEIGHT

BATCH

**BEST AVAILABLE COPY**

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed. Avoid contamination of food and foodstuffs. The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. Any exposed utensils or food handling surfaces should be thoroughly cleaned before reuse.

Remove pets, birds and cover fish aquaria before spraying.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.

Hold container upright, nozzle away from you and press actuator button spraying as directed.

**Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths:** Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

**Indian Meal Moths, Angoumois Grain Moths:** Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep areas closed for 15 minutes following treatment. Repeat application weekly or as needed.

**In Dairies and Milk Rooms:** For rapid control of house flies, mosquitoes, gnats, wasps and flying moths, close windows and doors and spray at a rate of one second per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**FAIRFIELD AMERICAN CORPORATION**

3932 Salt Road  
Medina, New York 14103

BYERONE - Registered trademark of Fairfield American Corporation

EPA Reg. No. 4816-523  
EPA Est. No. 279-NY-1

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
Harmful if swallowed. Do not apply over uncovered milk or food products.  
See rear panel for additional precautions.

BEST DOCUMENT AVAILABLE

NET WEIGHT

BATCH

ACCEPTED  
MAY 08 1979  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4816-523

may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.



Hold container upright, nozzle away from you and press actuator button spraying as directed.

Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths: Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

Indian Meal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep areas closed for 15 minutes following treatment. Repeat application weekly or as needed.

In Dairies and Milk Rooms: For rapid control of house flies, mosquitoes, gnats, wasps and flying moths, close windows and doors and spray at a rate of one second per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is complete.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



**FAIRFIELD AMERICAN CORPORATION**  
3932 Salt Road  
Medina, New York 14103

CODE 820.25

Name: PYRENONE<sup>®</sup> DAIRY AND FOOD PLANT AEROSOL  
Type: An essential use aerosol containing chlorofluorocarbon propellants  
Use Locations: Non-residential food handling areas, including dairies and milk rooms  
Insects Controlled: For use against flying insects only as an essential use product  
USDA Approval: Will be applied for in all areas of federally inspected meat and poultry plants  
Special Features:

- \* Self-propelled, ready-to-use
- \* Industrial Strength - low dosage required
- \* A true aerosol, no particle over 50 microns and 80% less than 30 microns
- \* Pyrenone gives rapid control of flying insects
- \* Pyrenone components have food additive status

Physical Data:

Specific Gravity @ 20°C:	1.24	Flame Extension:	8"
Delivery Rate @ 20°C:	1.53 g/sec	Drum Test:	Negative
P.S.I.G. @ 20°C:	55	Recommended Shelf Life:	3-4 years

Recommended Packaging: Can: 0.5 lb. three-piece tinplate or 35 oz. aluminum impact container  
Valve: Precision NN 0.018"/0.080"  
Actuator: Precision 0162-20 0.018

Package Sizes Registered:  
5, 8, 10, 12, 14 oz., 1 lb., 1 lb. 4 oz., 2 lb. 4 oz.

PYRENONE - Registered trademark of Fairfield American Corporation

4/17/79

0.20  
0.018



# MANZATE<sup>®</sup>

## FUNGICIDE

### ACTIVE INGREDIENT:

Maneb (Manganese ethylenebisdithiocarbamate) 80%  
(Manganese equivalent as metallic 16.5%)

### INERT INGREDIENTS

20%

EPA Reg No 352-173

Keep out of reach of children

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS

**CAUTION!** MAY IRRITATE EYES, NOSE, THROAT, AND SKIN  
MAY BE HARMFUL IF INHALED OR SWALLOWED

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks.

**FIRST AID** In case of contact flush skin or eyes with plenty of water for eyes, get medical attention.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

**IMPORTANT**—Never allow "Manzate" to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Manzate" as a fungicide and create vapors which may be flammable. Keep container closed when not in use.

## NET 3 LBS.

MADE BY DU PONT IN COLOMBIA

E. I. DU PONT DE NEMOURS & COMPANY (INC.)  
BIOCHEMICALS DEPARTMENT, WILMINGTON, DELAWARE

### GENERAL INFORMATION

Manzate is a fungicide which is effective against a wide range of fungi and molds. It is used for the control of fungi and molds on a variety of crops and materials. It is also used for the control of fungi and molds in the home and in other buildings. It is a contact fungicide and is effective against both active and dormant fungi and molds. It is a systemic fungicide and is effective against both active and dormant fungi and molds. It is a contact fungicide and is effective against both active and dormant fungi and molds. It is a systemic fungicide and is effective against both active and dormant fungi and molds.

### NOTICE OF WARRANTY

This product is warranted to be free from defects in material and workmanship. It is warranted to perform as described in the directions for use. It is not warranted to be effective against all fungi and molds. It is not warranted to be effective against all crops and materials. It is not warranted to be effective in all climates and conditions. It is not warranted to be effective in all cases. It is not warranted to be effective in all instances. It is not warranted to be effective in all situations. It is not warranted to be effective in all circumstances. It is not warranted to be effective in all cases. It is not warranted to be effective in all instances. It is not warranted to be effective in all situations. It is not warranted to be effective in all circumstances.

### DIRECTIONS FOR USE

**SPRAYS** Use Manzate as a spray for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials.

**DUSTS** Use Manzate as a dust for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials. It is used for the control of fungi and molds on a variety of crops and materials.

**ALMONDS** Use Manzate for the control of fungi and molds on almonds. It is used for the control of fungi and molds on almonds. It is used for the control of fungi and molds on almonds. It is used for the control of fungi and molds on almonds. It is used for the control of fungi and molds on almonds. It is used for the control of fungi and molds on almonds. It is used for the control of fungi and molds on almonds. It is used for the control of fungi and molds on almonds.

CONTINUED ON REAR PANEL

SEE REAR PANEL FOR MORE INFORMATION

ACTUAL SIZE