

CC L 820.75

PYRENECS® MULTI-PURPOSE INSECTICIDE

EPA Reg. No. 4816-525 EPA Est. No. 79-NY-1

Contains PYRENECS®

Professional Strength - Low Dosage Required

For use in Schools, Hospitals, Institutions, Supermarkets, Dairies, Hotels, Restaurants, Driveways, Motels, Drug Stores, Food Plants, Milk Rooms and Animal Quarters.

Provides Rapid Control of House Flies, Mosquitoes, Small Flying Moths (Millers), Gnats, Wasps and accessible stages of Cockroaches, Aits, Water Bugs, Silverfish, Flour Beetles, Weevils, Grain Mites and Spiders.

KILLS AND REPELS Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats.

Controls Garden Insects on Ornamental and Vegetable Plants.

Acceptable for Use to Day of Harvest on Edible Plants.

ACTIVE INGREDIENTS:

Pyrethrins	0.52
Piperonyl Butoxide, Technical	5.08
Petroleum Hydrocarbons	4.32
INERT INGREDIENTS	90.08
	100.00

Equivalent to 42 (butylcarbityl)(6-propylpiperonyl) ether and 12 related compounds.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See side panel for additional precautionary statements.

Pyrethrin - Registered Trademark of Fairfield
Pyrethrin - Registered Trademark of Fairfield

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

House Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths (Millers): Close all doors and windows and spray room at the rate of 2 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat as necessary.

Cockroaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes and Sow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Aits: Spray trails, nests and points of entry. Spray plants where possible. Repeat as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Carpet Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

In Animal Quarters and Milk Rooms: For rapid control of house flies, mosquitoes, gnats, wasps and small flying moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

SEE REVERSE SIDE

**FAIRFIELD
AMERICAN
CORPORATION**

3932 Salt Road
Manhasset Neck, New York 11030

To treat whole lots of beef cattle, dairy cattle and horses for attacks of stable flies, house flies, horn flies, eye flies, and other annoying insects, use the following method: Mix 3 seconds in 1 gallon of water. Spray carefully to spray back, with special care to the head. To protect from eye flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

Exposed Stages of Indian Meal Moths and Angoumois Grain Moths: Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Exposed Stages of Flour Beetles, Weevils and Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored food products in the home should be disposed of.

Control Garden Insects on Ornamental and Vegetable Plants: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

Acceptable for Use to Day of Harvest on Edible Plants

USE ON	TO CONTROL
Roses	Aphids, Rose Chafers, Japanese Beetles, Rose Slugs, Tent Caterpillars, Scales
Asters	Leafhoppers, Tarnish Plant Bugs, Lace Bugs, Thrips (exposed), Mealy Bugs
Danlias	Aphids, Red Spider Mites, Plant Bugs
African Violets	Broad Mites, Citrus Mealybugs
Azaleas	Azalea Lace Bugs, White Flies, Leaf Miners
Euonymus	Euonymus Scales, Aphids, Thrips (exposed)
Hydrangeas	Aphids, Red Spider Mites, Hydrangea Leafhoppers

USE ON	TO CONTROL
Cucumbers	Cucumber Beetles, Red Spider Mites, Aphids
Peppers	Leaf Miners, Leaf Rollers, Fleas
Peppers, Eggplants, Potatoes	Colorado Potato Beetles, Leafhoppers, Blister Beetles, Flea Beetles, Green Peach Aphids
Cucumbers, Squash, Melons	Colorado Potato Beetles, Leafhoppers, Blister Beetles, Flea Beetles, Green Peach Aphids
Beans	Colorado Potato Beetles, Leafhoppers, Blister Beetles, Flea Beetles, Green Peach Aphids
Sweet Corn	Colorado Potato Beetles, Leafhoppers, Blister Beetles, Flea Beetles, Green Peach Aphids

STORAGE AND DISPOSAL

Storage: Store in a cool, dry area away from heat or open flame.

Disposal: Wrap container and dispose of in trash. Do not incinerate.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. All food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by disposal of waste or cleaning of equipment.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

Buyer assumes all risks of storage or handling of this product not in strict accordance with directions given herewith.