

BEST AVAILABLE COPY

Reg. # 4816-525
CODE 820.75

Pyrenone® MULTI-PURPOSE INSECTICIDE

EPA Reg. No. 4816-525

EPA Est. No. 279-NY-1

CONTAINS PYRENONE®

PROFESSIONAL STRENGTH - LOW DOSAGE REQUIRED

For use in SCHOOLS, HOSPITALS, INSTITUTIONS, SUPERMARKETS, DINING ROOMS, RESTAURANTS, DRIVE-INS, MOTELS, DRUG STORES, FOOD PLANTS, MILK ROOMS and ANIMAL QUARTERS

Provides rapid control of HOUSE FLIES, MOSQUITOES, SMALL FLYING MOTHS (Millers), GNATS, WASPS and accessible stages of COCKROACHES, ANTS, WATER BUGS, SILVERFISH, FLOUR BEETLES, WEEVILS, GRAIN MITES and TERMITES

KILLS AND REPELS STABLE FLIES, HORSE FLIES, FACE FLIES, DEER FLIES, HOUSE FLIES, MOSQUITOES and GNATS

Controls garden insects on Ornamental and Vegetable Plants

AVAILABLE FOR USE TO DAY OF HARVEST ON EDIBLE PLANTS

ACTIVE INGREDIENTS

| | |
|------------------------------|--------|
| Pyrenone | 0.5% |
| Pyrenone Butoxide, Technical | 5.0% |
| INERT INGREDIENTS | 94.5% |
| | 100.0% |

*Equivalent to 47 (4-ethylarbityl)(6-propylpiperonyl) ether and 12 related compounds

© 1975, registered trademark of Fairfield American Corporation

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See rear panel for additional precautions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

House Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths (Millers): Close all doors and windows and spray room at the rate of 2 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat as necessary.

Cockroaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes and Sow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor coverings where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Carpet Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

IN ANIMAL QUARTERS and MILK ROOMS: For rapid control of House Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

TO TREAT ANIMALS: To protect beef cattle, dairy cattle and horses from attacks of Stable Flies, Horse Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats, spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.



Exposed Stages of Indian Meal Moths and Angoumois Grain Moths: Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for that purpose.

Exposed Stages of Flour Beetles, Weevils and Grain Mites: Spray into cracks, and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored products in the home should be disposed of.

SEE REVERSE SIDE

NET CONTENTS _____

FAIRFIELD 784



**FAIRFIELD
AMERICAN
CORPORATION**
238 WILSON AVENUE
NEWARK, N.J. 07105

CODE 820.75

Control Garden Insects on Ornamental Vegetable Plants: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

Acceptable for Use To Day of Harvest on Edible Plants.

Use on Ornamentals (Roses, Asters, Dahlias, African Violets, Azaleas, Euonymus, Magnolias, Junipers, Lilacs) to control Aphids, Rose Chafer, Japanese Beetles, Leafhoppers, Tomato Hornworms, Scalps (including Euonymus and Juniper Scales), Spider Mites, Japanese Stink Bugs, Lace Bugs (including Azalea Lace Bugs), Thrips, Whiteflies, Mealybugs (including Citrus Mealybugs), Mites (including Red Spider, Spider and Red Spider), Plant Bugs, White Flies, Leaf Miners, Hydrangea Leaf-Miner, Hydrangea and Leaf Killers.

Use on Vegetables (Peppers, Eggplants, Potatoes, Cucumbers, Squash, Melons, Beans, Sweet Corn) and Tomatoes to control Flea Beetles, Hornworms, Stink Bugs, Cucumber Potato Beetles, Aphids (including Green Peach Aphids), Leafhoppers, Spider Beetles, Striped Cucumber Beetles, Mexican Bean Beetles, White Flies, Mealybugs and Japanese Beetles.

STORAGE AND DISPOSAL

Store in a cool, dry area away from heat or open flame. Do not transport or store below 32°F.

Seal container in newspaper and dispose of in trash.

PRECAUTIONARY STATEMENTS RELATING TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash all equipment, benches, walls, etc., where exposed food will be handled, with potable water. All food processing surfaces and utensils should be covered during treatment and thoroughly washed before use.

Remove pets, birds, and other household animals before spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching finger to back of throat. Never give anything by mouth to an unconscious or convulsing person.
IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.
IF IN EYES OR ON SKIN: Flush affected areas with plenty of water. Contact a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by disposal of wastes or cleaning of equipment.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

10-15-04

10-15-04