

CODE 721.90

PYRENONE® Food Plant Fogging Insecticide

EPA REG NO 4816-534

NET CONTENTS 1 QUART
NET CONTENTS 1.65 LBS.

A Concentrated PYRENONE Product

Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
Petroleum Distillate	94.5%
	<u>100.0%</u>

*Equivalent to 4% (butylcarbityl)(6-propiperonyl) ether and to 1% related compounds

PYRENONE - Fairfield American trademark

CAUTION
KEEP OUT OF REACH OF CHILDREN.

72190 1G 277

CAUTION: Harmful if swallowed. Avoid contamination of feed and foodstuffs.

Remove pets, birds and cover fish aquaria before spraying. This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

DIRECTIONS

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths:

Close windows and doors and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. In Official Meat and Poultry Establishments use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. In other than Official Meat and Poultry Establishments a combination of spot surface treatment and fogging is recommended.

See enclosed sheet for directions for use

BATCH



FAIRFIELD AMERICAN
CORPORATION
100 NIAGARA ST.
MIDDLEPORT, N.Y. 14105

4816-534

03/31/1977

1/2

DIRECTIONS

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths:

Close windows and doors and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. In Official Meat and Poultry Establishments use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. In other than Official Meat and Poultry Establishments a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worm, grain mites, and caddis: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.



**FAIRFIELD
AMERICAN**
CORPORATION
100 NIAGARA ST.
MIDDLEPORT, N.Y. 14105

CODE 721.90

PYRENONE® Food Plant Fogging Insecticide

EPA REG NO 4816-534

A Concentrated PYRENONE Product

Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
Petroleum Distillate	94.5%
	100.0%

*Equivalent to 4% (butylcarbityl)(6-propiperonyl) ether and to 1% related compounds

PYRENONE - Fairfield American trademark

CAUTION

Keep Out Of Reach Of Children

CAUTION: Harmful if swallowed. Avoid contamination of feed and foodstuffs.

Remove pets, birds and cover fish aquaria before spraying. This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

72190 C

8/2