

CODE 72190 ACCEPTED

Pyrenone[®] FOOD PLANT FOGGING INSECTICIDE

MAY 8 1986
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 72190-54

EPA Reg. No. 4816-534 EPA Est. No. 279-NY-1
Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist

ACTIVE INGREDIENTS:

Pyrethrins.....	0.5%
*Piperonyl Butoxide, Technical.....	5.0%
FINEST INGREDIENTS:	94.5%
	100.0%

*Equivalent to 4.0% (butylcarbityl)(6-propylpiperonyl) ether and 1.0% related compounds.

†Contains petroleum distillate

PYRENONE - Registered Trademark of Fairfield American Corporation.

KEEP OUT OF REACH OF CHILDREN
CAUTION
See rear panel for additional precautions

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SPACE APPLICATIONS:


For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths.

Close windows and doors and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist of fog. In Official Meat and Poultry Establishments use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes.


Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. In other than Official Meat and Poultry Establishments a combination of spot surface treatment and fogging is recommended.

IN BARN, POULTRY HOUSES and STORAGE ROOMS

To control house flies, stable flies and horn flies in Barns, Poultry Houses, Warehouses, Storage Bins and Storage Rooms: dilute 1 part of material with 4 parts of oil. Apply as a fine mist over the livestock in stanchions or holding pens. Use a fogger or mist generator to direct spray above the animals and to the center of the area where animals are housed. Apply 1 ounce per 1000 cubic feet uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in Dairy Barns, aerate the barn thoroughly before milking. Repeat daily in the morning and afternoon, or as required.



NET CONTENTS:



72190 286

SURFACE SPRAYING

To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, mealworm, grain mites, and caddis: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces apply at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make a localized application to floor and baseboards: Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as necessary.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining. Store clothes in moth tight containers or repeat treatments monthly.

OUTDOOR APPLICATIONS:

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy areas, stagnant pools or ponds and shorelines. Apply at a rate of 10 fluid ounces per acre (0.2 fluid ounce per 1000 square feet).

Using truck-mounted conventional misting equipment traveling at 5 m p h. to treat a 300 swath, calibrate equipment to deliver 0.23 fluid ounce per minute.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.


CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by an approved waste disposal facility.

CONTAINERS ONE GALLON AND SMALLER: Store in original container in a cool dry place inaccessible to children and pets. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

SEE REVERSE SIDE



FAIRFIELD AMERICAN CORPORATION
238 WILSON AVENUE
NEWARK, N.J. 07105



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs. The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment and thoroughly washed with an effective cleaning compound followed by a potable water rinse before using. Remove pets, birds, and cover fish aquaria before spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center immediately. Do NOT induce vomiting unless directed to do so by a physician. Vomiting may cause aspiration pneumonia.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.


IF IN EYES OR ON SKIN: Flush affected areas with plenty of water. Contact a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BATCH



Printed in U S A