

# GREENHOUSE SYNTHRIN® WHITEFLY SPRAY CONCENTRATE

NET WEIGHT

2,2,4-trichloro-1,1,2-dimethyl-3-(2,4,6-trichlorophenyl)butane-3-carboxylate	24.30
Alkylamine	3.30
Surfactant	34.40
Inert petroleum hydrocarbons	33.00
<b>NET INGREDIENTS</b>	<b>5.00</b>
	<b>100.00</b>

Contains 100% active ingredient. 30 (±) oz. and 100 (±) g. net wt. U.S. Patent No. 2,465,007

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS - CHEMICAL**  
Harmful if swallowed. Irritant to skin. Avoid breathing spray mist. Avoid contact with eyes. Do not spray on food or feed crops.

**FIRST AID**  
In case of contact with eyes, immediately flush eyes with plenty of water for 15 minutes. For more, call a physician.

**ENVIRONMENTAL HAZARDS**  
This product is toxic to fish and birds. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**  
Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**  
Do not reuse empty container. Wrap container and put in trash collection.

**CONTACT SPRAY** - For whiteflies, apply at 1 pint per 100 gallons of water or 1 teaspoonful per gallon. Spray upper and lower surface of plant foliage to point of runoff. Spray in late afternoon and at night only, when the temperature ranges from 50°F to 72°F. Repeat application weekly for four weeks or longer for control of adults and nymphs. Complete spray control on poinsettias before bloom form.

where fewer applications are preferred, spray at the rate of 1 quart per 100 gallons of water or 2 teaspoonfuls per gallon. Repeat applications. Apply spray as vented or above.

**FACE APPLICATIONS** - Do not allow plants to be treated for whitefly control in a greenhouse. First mist toward upper corners of the greenhouse, then apply evenly throughout the area. Keep nozzle at least three feet from plant foliage. Apply at the rate of 1 fl. oz. four quarts mixed in 3 fl. oz. of methylene chloride per 10,000 cu. ft. This is equivalent to 1 gram active ingredient per 1000 cu. ft. Keep treated area closed for at least four hours. Do not walk in treated area. Ventilate the area after treatment is completed. Apply at weekly intervals for at least four weeks. Applications should be made during the late afternoon or at night when the temperature range is 50°F to 72°F.

**GREENHOUSE PESTS**: For infestations of whiteflies on begonias, fuchsia, salvia, poinsettia, chrysanthemums, coleus, geraniums, ageratum, petunia, zinnia, calendula, marigolds, roses, azaleas and ...

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**Keep Out Of Reach Of Children**  
**WARNING**  
See side panels for additional precautionary statements.

SYNTHRIN - Registered trademark of Fairfield American Corporation  
EPA Reg. No. 4816-545  
EPA Est. No. 279-NY-1

NET CONTENTS \_\_\_\_\_ GALLONS  
NET CONTENTS \_\_\_\_\_ POUNDS  
NET CONTENTS \_\_\_\_\_ LITERS  
NET CONTENTS \_\_\_\_\_ KILOS

**ACCEPTED**  
**06 JUL 1979**  
United States Dept. of Agriculture  
EPA, and the Federal Insecticide, Fungicide, and Rodenticide Act,  
as amended, in the pesticide  
registered under  
EPA Reg. No. 4816-545

BATCH \_\_\_\_\_

