Ready-to-use for control of residential, indusreial and parden insect pasts - Non-food areas of industrial buildings.

Tontains a combination of two highly active synthetic Pyrethroid Compounds

ACTIVE INGREDIENTS


MOTITE HORIEDICITIO	
*Totramethrin	0.250%
Related Compounds	
**Resmethrin	
Related Compounds	0.046%
"INERT INGREDIENTS	99,420%
	100.000%

- *1-cyclohexene-1,2-dicarboximidomethyl 2,2-dimethyl-3 (2methylpropenyl) cyclopropanecarboxylate
- **Cis/trans isomer content: Max 43% (+) cis and min. 70% (+) trans.
- ***Contains aromatic petroleum hydrocarbons and petroleum distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautions.

EPA Peg. No. 4816-547 EP4 LSC. No. 279-NY-1



NET CONTENTS____

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Food and drink should be removed or covered during treatment. Avoid breathing of vapor or contact with skin, eyes or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get Imedical attention if irritation persists. Do not smoke while spraying. Remove pets, birds and cover fish aquariums before spraying. Do not use on edible crops. Do not use where food is commercially prepared. Do not use in serving areas while food is exposed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Shake well before using.

INDOORS:

FOR CONTROL OF RESIDENTIAL AND INDUSTRIAL INSECTS:

> Flies Mosquitoes

Gnats.

Small Flying Moths

Close doors and windows and shut off ventilating systems. Using a conventional mechanical or compressed air spraying or fogging equipment, apply as a fog or fine mist, directing the spray toward the upper portions of the room and toward · windows and other light sources which attract these insects, thoroughly fill the area with mist using 1 to 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Vacate after treatment and ventilate before reoccupying. Repeat treatment as needed.

Hornets.

Direct the spray at these insects aften possible a Spray thoroughly Thto nests, hading and brending placed when inneces have selled at neglet.

Spray trails, nests and prints of entry. Spray directly on ants when possible. Repeat at necessary.

> Cockroaches Sowbuas Boxelder Bugs Centipedes

Crickets Earwios Grain Hites Spiders

Apply as a coarse droplet spray directed into cracks, crevices and other hiding places along baseboards, around the base of machinery and under pallets. Spray thoroughly contacting insects directly whenever possible. Spray badly infested areas daily. In other areas, repeat treatments twice weekly or when reinfestation occurs.

INDOORS AND OUTDOORS:

HOUSE PLANTS AND GREENHOUSES AND ORNAMENTALS OUTDOORS

Petunia

Snapdragon

Rose

Azalea

Euonymus

For control of: Aphids

Two-spotted spider mites

Plants:

lvy

African Violets Aralia

Lantana Heathe bella paim

Christmas Cherry

Chrysanthenum Cineraria


Diffenbachia Euschia

Gloxinia Marigold Zinnia

Di ect the spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

BEST AVAILABLE COPY

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.



高端を到りはほりは CORPORATION

3932 Salt Road Matien to Ver term