

MULTI-PURPOSE RESIDUAL 70 W.P.

4816-550 PM 12 1002
 CODE 123.45

Contains PROPOXUR

EPA REG. No. 4816-550 EPA EST. No. 4972-FL-1


To be applied only by or under the supervision of Trained Personnel responsible for Insect Control Programs.

An insecticide for use in and around homes, apartment buildings, stores, warehouses and animal research facilities.

This jar contains 5x2 ounce water-soluble pouches of Multi-Purpose Residual 70 W.P. which dissolve and mix readily in water to form a suspension. Apply with most hand or power operated sprayers.

ACTIVE INGREDIENTS

*2-(1-Methylethoxy) phenol methyl-
 carbamate 70%
 INERT INGREDIENTS 30%



KEEP OUT OF REACH OF CHILDREN
WARNING
 See back panel for additional precautions.

NET CONTENTS: 10 POUCHES

BATCH: 1234567890

Printed in U.S.A.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Avoid excessive handling of pouches or exposure to moisture.

For control of the following insects in and around Homes, Apartment Buildings, Stores and Warehouses.

INDOORS

- | | |
|--|------------|
| Ants | Silverfish |
| Cockroaches | Spiders |
| Fleas | Ticks |
| Millipedes | Waterbugs |
| Sawtoothed Grain Beetles (exposed stages). | |

Dissolve 1 pouch in 1 gallon of water to make a 1 1% concentration. Apply as a residual spray to baseboards, window frames, the undersides of shelves and drawers, under sinks and stoves and to other places where insects hide. Also spray cracks, crevices and surfaces where insects may crawl when they come out of hiding. Continue applications as needed with a maintenance application using a 0.5% solution (1 pouch in 2 gallons of water). Do not spray foodstuffs or house plants. Do not use as a space spray. Do not treat animals. Treat pets with an E.P.A. approved product registered for use on animals.

Pest Control in Meal Packing and Food Processing Plants, Food Storage Areas, Stores, Warehouses, Restaurants, and other Food Handling Establishments (places other than private residences in which exposed food is held, processed, prepared or served.) To obtain initial control of ants, cockroaches, millipedes, silverfish, spiders, waterbugs, and to aid in the control of sawtoothed grain beetle (exposed stage), use 1 pouch in 1 gallon of water to make a 1 1% concentration. Continue applications as needed using a maintenance rate of 0.5% (1 pouch in 2 gallons of water).

Non-Food Areas: Apply to floors, walls, ceilings, or other infested areas.

Food Areas: Limited to Crack and Crevice Treatment Only - Apply a small amount of material directly into cracks and crevices such as expansion

joints between different elements of construction or between equipment bases and the floor, wall voids, or hollow equipment legs where the insects noted above hide. Do not use this product in conduits, motor housings, or electrical switch boxes. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

Applications of this product in the food areas of food handling establishments, other than as a crack and crevice treatment, are not permitted.

Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestations. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

For control of the following pests in Research Animal Facilities:


- | | |
|---|------------|
| Ants | Silverfish |
| Cockroaches | Spiders |
| Fleas | Ticks |
| Millipedes | Waterbugs |
| Sawtoothed Grain Beetles (exposed stages) | |

Follow the above directions. The use of Baygon in and around research animal's quarters for pest control will not induce changes in liver microsomal enzyme activity in these animals. Multi-Purpose Residual 70 W.P. should NOT be applied in portions of the buildings where there is direct contact by the animals. Avoid contamination of food. Do not use as a space spray. Do not treat animals. Treat animals with an E.P.A. approved product registered for use on animals.

OUTDOORS

- | | |
|--------------|------------|
| Ants | Millipedes |
| Cockroaches | Mosquitoes |
| Clover Mites | Sowbugs |
| Crickets | Spiders |
| Fleas | Earwigs |
| Ticks | Waterbugs |

Dissolve 1 pouch to 1 gallon of water to make a 1 1% concentration. Apply as a residual spray or with a paint brush to surfaces of buildings, porches, screens, window frames, patios and garages where these insects congregate. Repeat as necessary. Do not treat plants (lawns, flowering plants, shrubs, trees, etc) with this concentrate.



FAIRFIELD AMERICAN

201 Rt 17 North
 Rutherford NJ 07070

STORAGE AND DISPOSAL

To Control Sand Flies or Punkles:

Dissolve 1 pouch in 1 quart water to make a 4 2% concentration. Apply with a paint brush to outside surfaces of door and window frames and other screened areas where these insects may enter buildings. Thorough coverage is necessary for maximum control.

For Control of Sodworms (lawn moths) on Lawns and Turfs:

Dissolve 1 pouch in 10 to 15 gallons of water to make a 0.11% to 0.07% concentration. Apply to 700 square feet of area. For best results, mow the lawn and rake dead leaves from damaged spots, water the lawn thoroughly and allow grass to dry before spraying. If possible do not water again for several days. Three applications at monthly intervals may be required for maximum control. In most areas, the first application should be made in May or June. Consult your local State Agricultural Experiment Station or Extension Service for more specific information regarding timing of application.

For control of Chinch Bugs, Hunting Bill bugs, Ticks, Chiggers, and Bluegrass Bill-bugs on Lawns and Turfs:

Apply 2 pouches per 1000 square feet of area. Do not use less than 15 gallons of water per 1000 square feet. For best results use 30 to 40 gallons of water per 1000 square feet. Repeat as necessary. For most effective control of chinch bugs, ticks and chiggers, apply during the spring and summer months, or as recommended by your State Agricultural Extension Service. For best results, lawns should be watered before applying Baygon.

Low Volume Spray for Control of Adult Mosquitoes:

Dissolve 1 to 2 pouches in one gallon of water (0.05 to 0.175 lb. active) per acre for ground or aerial application.

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Unused pouches should remain in jar. Do not store this jar under wet conditions. Handle carefully when stored at temperature of less than 50°F to avoid breakage of soluble pouch.

DISPOSAL: Do not reuse empty jar. Wrap and put in trash collection. Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS WARNING

May be fatal if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin or on clothing. In case of prolonged exposure, wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and warm water. Do not store near feed or food products. Do not contaminate food. Wash hands, arms and face thoroughly with soap and water before eating or drinking. Wash all contaminated clothing with soap and hot water before reuse. Since this is a water suspension material, do not apply to rugs, carpets, draperies, wallpaper or similar materials that may be stained or watermarked. Keep children and pets off treated areas until surfaces are dry. Do not apply to animals or humans. Do not use in animal barns or poultry houses for fly control.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes or on Skin: Flush with plenty of water and get immediate medical attention.

If Swallowed: Drink a glass of water and induce vomiting by touching finger to back of throat. Get medical attention.

If Inhaled: Remove victim to fresh air. Apply artificial respiration and oxygen if indicated.

NOTE TO PHYSICIAN Atropine Sulfate is antidotal

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife. Birds feeding on treated areas may be killed. Do not apply directly to lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.