

- Contact Spray - Diluted for Use with Sprayers that Give Large Wetting Spray Particles to Kill
Cockroaches and Accessible Stages of Granary
Insects: Granary Weevils, Rice Weevils, Confused
Flour Beetles, Red Flour Beetles, Meal Moth Lar-
vae and Adults, Caddis, Grain Mites and Saw-
toothed Grain Beetles: Remove or cover exposed
food and close room as described above. Dilute
concentrate with oil at the rate of 1 part con-
centrate plus 29 parts of oil or 4½ ounces per
gallon and mix well. Spray outside walls of
grain bins, flooring, about base of milling ma-
chinery, walls and baseboards, areas behind cab-
inets, around pipes and other insect harborages
using 1 gallon spray per 1000 square feet of
surface. Force spray into all cracks and crev-
ices. Repeat applications as necessary.

Contact and Space Spray - Undiluted - To Kill
Cockroaches, Silverfish, Earwigs, Ants, and
Spiders in Sheltered Situations: With ULV spray-
ing device, apply the spray into all cracks and
crevices in woodwork, walls, floors, underneath
sinks, behind pipes and in all places where
these insects find shelter. Contact as many in-
sects as possible. Do not allow spray to contact
stored food. Use at a dosage rate of approxima-
tely ½ fluid ounce per 1000 cubic feet of room
space. Repeat dosage as necessary for control.
Leave the room closed for one hour. Do not re-
main in treated areas and ventilate the area
before re-entry. Do not apply this undiluted
spray in facilities operating under USDA poul-
try, meat, rabbit and egg products inspection
programs.

Buyer assumes all risks of use, storage or
handling of this material not in strict ac-
cordance with directions given herewith.

NET CONTENTS _____ GALLONS
NET CONTENTS _____ POUNDS
NET CONTENTS _____ LITERS
NET CONTENTS _____ KILOS

BATCH

