

Reg. # 4816-560

PM-17

1971

# DE 923.65 PYRANHA™ FLYING INSECTICIDE

Not Reviewed. Registrant claims to be in accordance with Registration of Pesticide Product Guidance for Phase 2 response. Page 2.12, Active to Inert Change in Status.

Controls house flies, stable flies, horse flies, fruit flies, mosquitoes, gnats, and small flying moths. Also controls bedbugs and mites in poultry houses when applied as directed.

For use in horse barns, kennels, dairy barns, milking parlors, milk rooms, poultry houses, food processing plants, bottling plants, canneries, flour mills and grain elevators.

### ACTIVE INGREDIENTS

Pyrethrins.....	1.82%
Piperonyl Butoxide, Technical.....	18.94%
**INERT INGREDIENTS.....	79.24%
	<u>100.00%</u>

<sup>1</sup>Equivalent to 15.15% (butylcarbityl)(6-propyl-piperonyl) ether and 3.79% related compounds

\*\*Contains Petroleum Distillate  
CHEM-I-MATIC, PYRANHA & STABLE MASTER - Trademarks of Chem-I-Matic, Inc.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side panel for additional precautions.

EPA Reg. No. 4816-560  
EPA Est. No. 279-NY-1

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

CAUTION: Harmful if swallowed. Avoid contamination of feed and foodstuffs. The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be covered or removed during treatment or thoroughly cleaned before using.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely. Do not contaminate water by cleaning, rinsing or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

For use exclusively in the formulation of Pyranha insecticides, to be dispensed through Chem-I-Matic automatically timed Stablemaster spraying systems.

Pyranha Flying Insecticides is designed to be diluted in water just prior to use. To formulate a spray containing 0.1% pyrethrins and 1.0% piperonyl butoxide, use 3 gallons with 51 gallons of water or 7 fluid ounces of concentrate per gallon. The Pyranha should be added to the water slowly, while stirring or agitating. Holding tanks should be emptied and thoroughly cleaned at frequent intervals to prevent biological growth.

When used in Stablemaster Automatic Spray Systems, add the diluted spray to storage tank. Re-seal tank. Set nozzles to deliver 1 ounce of liquid per minute.

Set timer to operate 2 three-minute cycles daily so that each nozzle delivers a total of 6 ounces of diluted spray each 24 hours. Properly locate each nozzle to cover a maximum of 2000 cubic feet of space.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

NET CONTENTS _____	GALLONS
NET CONTENTS _____	LITERS
NET CONTENTS _____	POUNDS
NET CONTENTS _____	KILOS

**BATCH**

**BEST AVAILABLE COPY**

