

CODF 923 55 PYRANHATM INSECTICIDE WITH REPELLENT


KEEP OUT OF REACH OF CHILDREN

CAUTION


BATCH

4316-270

DATE RECEIVED _____
BY _____
DATE RECEIVED _____
BY _____
DATE RECEIVED _____
BY _____



**FAIRFIELD
AMERICAN
CORPORATION**
3902 SALT ROAD
MADISON, N.J. 07042



CODE 923.55 PYRANHA INSECTICIDE WITH REPELLENT

Reg. # 4816-563

PM-17

191

Controls house flies, horn flies, stable flies, horse flies, mosquitoes and gnats

For use in horse barns, kennels and other non-meat or dairy animal quarters. Not for use on beef or dairy animals, poultry or in farm structures or buildings housing food producing animals or poultry or milk rooms.

ACTIVE INGREDIENTS

*Pyrethrins	1.37%
*Piperonyl Butoxide, Technical	13.71%
Butoxypropylene Glycol	27.56%
**INERT INGREDIENTS	57.36%
	<u>100.00%</u>

¹Equivalent to 10.97% (butylcarbityl)(6-propyl-piperonyl) ether and 2.74% related compounds ;

**Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional precautions.

EPA Reg. No. 4816-563
EPA Est. No. 279-NY-1

CHEM-I-MATIC, PYRANHA & STABLE MASTER -
Trademarks of Chem-I-Matic Inc.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

CAUTION: Avoid breathing of spray. Avoid contamination of food and feedstuffs. Harmful if swallowed. Do not use on animals intended for food.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

BEST AVAILABLE COPY

NOV 1 1979

4816-563

NET CONTENTS _____ GALLONS
NET CONTENTS _____ LITERS
NET CONTENTS _____ POUNDS
NET CONTENTS _____ KILOS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Pyranha Insecticide is designed to be diluted in water just prior to use. To formulate a spray containing 0.1% pyrethrins, 1.0% piperonyl butoxide and 2.0% Stabilene, use 4 gallons in 49 gallons of water for 10 fluid ounces of concentrate per gallon. Pyranha Concentrate should be added to the water, slowly, while stirring or agitating. Holding tanks should be emptied and thoroughly cleaned at frequent intervals to prevent biological growth.

Add the diluted spray to storage tank of Stable Master[®] Automatic Space Spray system. Reseal tank. Set nozzle to deliver 1 ounce per diluted spray per minute.


Set timer to operate 2 three-minute cycles daily so that each nozzle delivers a total of 6 ounces of diluted spray each 24 hours.

Properly locate each nozzle to cover a maximum of 2,000 cubic feet of space.

For control of bedbugs and mites in poultry houses, spray crevices of roost holds, cracks in walls and nests and other areas where bedbugs and mites might hide. Repeat treatment as needed.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

BATCH



**FAIRFIELD
AMERICAN
CORPORATION**

3rd Hill Road
Madira, New York 14103