

BEST DOCUMENT AVAILABLE

Pyrethron® GARDEN SPRAY

PRESSURIZED

CODE

An Effective, Fast Acting Insecticide For Use On Vegetables and Ornamental Plants

Approved for Use To Day of Harvest on Edible Plants

ACTIVE INGREDIENTS

Permethrin 0.5%
Piperonyl Butoxide 1.0%
Pyrethrin 0.5%
Piperonyl Butoxide 1.0%
INERT INGREDIENTS 98.0%

*Equivalent to 2.4% permethrin in appropriate formulations.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional product usage labeling.

PRESENTED BY: Fairfield American Corporation
A Division of The Clorox Company

EPA Reg. No. 40-147
EPA Est. No. 27-NM-1

PRECAUTIONARY STATEMENT HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

Keep out of reach of children. Avoid breathing of spray or contact with eyes or skin. Wash thoroughly after use. Do not contaminate feed and foodstuffs.

ENVIRONMENTAL HAZARD

This product is toxic to fish. Do not use near fish ponds or streams. Do not use near streams or ponds.

MEDICAL HAZARD

Contents under pressure. Do not puncture. Do not store near heat or open flame. Do not store below freezing temperatures. Exposure to temperatures above 130°F may cause bursting. Do not throw container into fire or incinerator.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

DISPOSAL: Wrap container and dispose of in trash. Do not incinerate.

Use can well before using.

This insecticide may be used outdoors in gardens or in greenhouses throughout the growing season or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

For best results, use throughout the growing season to prevent buildups of insect infestations. Spray plants from a distance of 1 1/2 to 2 feet using short bursts of 1 second. To prevent injury to plants, apply visibly wetting the foliage.

As soon as the early morning or late evening when the air is still, treat up all foliage with particular attention to the undersides of leaves. Contact insects directly whenever possible. Repeat treatment every 7-10 days or as reinfestation occurs.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatments as needed to control infestations. May be applied on day of harvest.

NET CONTENTS _____ GALLONS
NET CONTENTS _____ LITERS
NET CONTENTS _____ POUNDS
NET CONTENTS _____ KILOS

USE ON	TO CONTROL
Asparagus	Aphids, Leafhopper
Beans	Aphids, Leafhopper, White Fly, Mexican Bean Beetle, 12-Spotted Cucumber Beetle
Broccoli, Cabbage, Brussels Sprouts, Cauliflower	Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug
Celery	Green Peach Aphid, Cabbage Looper, Leafhopper
Cranberries	Fireflies, Leafhoppers
Cucumbers, Melons, Squash	Flea Beetles, Green Peach Aphid
Tomatoes	Blister Beetles, Colorado Potato Beetles, Flea Beetles, Green Peach Aphids, Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm
Mustard Greens, Kale, Collards, Turnips	Imported Cabbageworm, Diamondback Caterpillars, Aphids, Cabbage Looper
Peppers	Green Peach Aphids
Potatoes	Aphids, Colorado Potato Beetle, Flea Beetles, Leafhoppers
Radishes	Aphids, Flea Beetles
Spinach	Aphids, Cabbage Looper, Leafhoppers
Sweet Corn	Japanese Beetles
Tomatoes	Flea Beetles, Colorado Potato Beetles, Stink Bugs, Green Peach Aphids

FOR USE ON ORNAMENTALS: African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, daisies, dogwood, geraniums, gladioli, marigolds, roses, rubber plants. Wandering Jew: Insects to be controlled are aphids, flea beetles and leafhoppers.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

BATCH

ACCEPTED
4816-564



FAIRFIELD AMERICAN CORPORATION
3332 Salt Road
Medina, New York 14103