

CODE

Pyrenone® GARDEN SPRAY

PRESSURIZED

An Effective Fast Acting Insecticide For Use On Vegetable and Ornamental Plants

Accepted for Use To Day of Harvest on Edible Plants

ACTIVE INGREDIENTS:

Pyrethrins.....	0.050%
*Piperonyl Butoxide, Technical.....	0.500%
Petroleum Distillate.....	8.075%
INERT INGREDIENTS:	91.375%
	100.000%

*Equivalent to 0.40% (Butylcarbityl)(6-propyl-piperonyl) ether and 0.10% related compounds

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional precautionary labeling.

•PYRENONE- Registered trademark of Fairfield American Corporation

EPA Reg. No. 4816-564
EPA Est. No. 279-NY-1

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Keep out of reach of children. Avoid breathing of spray or contact with eyes or skin. Wash thoroughly after use. Do not contaminate feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not use near fishbowls or aquaria. Keep out of lakes, streams or ponds.

PHYSICAL HAZARDS

Contents under pressure. Do not puncture. Do not store near heat or open flame. Do not store below freezing temperatures. Exposure to temperatures above 130°F may cause bursting. Do not throw container into fire or incinerator.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

DISPOSAL: Wrap container and dispose of in trash. Do not incinerate.

Shake can well before using.

This insecticide may be used outdoors in gardens or in greenhouses throughout the growing season or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

For best results, use throughout the growing season to prevent buildups of insect infestations. Spray plants from a distance of 1½ to 2 feet using short bursts of 1 second. To prevent injury to plants, avoid visibly wetting the foliage.

Apply in the early morning or late evening when the air is still, treating all foliage with particular attention to the undersides of leaves. Contact insects directly whenever possible. Repeat treatment every 7-10 days or as reinfestation occurs.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatments as needed to control infestations. May be applied on day of harvest.

NET CONTENTS _____	GALLONS
NET CONTENTS _____	LITERS
NET CONTENTS _____	POUNDS
NET CONTENTS _____	KILOS

USE ON	TO CONTROL
Asparagus	Asparagus Beetle
Beans	Aphids, Leafhoppers, White fly, Mexican Bean Beetles, 12-Spotted Cucumber Beetle
Broccoli, Cabbage, Brussels Sprouts, Cauliflower	Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug
Celery	Green Peach Aphid, Cabbage Looper, Leaf-tier
Cranberries	Fireworms, Leafhoppers
Cucumbers, Melons, Squash	Flea Beetles, Green Peach Aphids
Eggplants	Blister Beetles, Colorado Potato Beetles, Flea Beetles, Green Peach Aphids
Lettuce	Cabbage Looper, Green Peach Aphids, Diamondback Moth Larvae, Imported Cabbageworms
Mustard Greens, Kale, Collards, Turnips	Imported Cabbageworms, Diamondback Caterpillars, Aphids, Cabbage Looper
Peppers	Green Peach Aphids
Potatoes	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers
Radishes	Aphids, Flea Beetles
Spinach	Aphids, Cabbage Looper, Webworms
Sweet Corn	Japanese Beetles
Tomatoes	Flea Beetles, Colorado Potato Beetles, Stink Bugs, Green Peach Aphids

FOR USE ON ORNAMENTALS (African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, dogwood, geraniums, gladioli, marigolds, roses, rubber plants, Wandering Jew): Insects to be controlled are aphids, flea beetles and leafhoppers.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

BATCH

ACCEPTED
NOV 9 1979
5164
4816-564



FAIRFIELD AMERICAN CORPORATION
3932 Salt Road
Medina, New York 14103

4816-564
11/06/1979