

Cover or tent or exposed food and cover food handling surfaces in edible product areas of food handling establishments. Close doors and shut off air conditioning or ventilating equipment. Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal one gallon of diluted spray. Apply at the rate of 1-2 fluid ounces per 1000 cubic feet, filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment if reinfestation occurs.

STALL QUARTER USE (Cattle barns, horse barns, poultry houses, swine houses, zoos): As a space spray, diluted for use in conventional chemical fogging or spraying equipment.

Control: Flies, Mosquitoes, Small flying moths, Gnats
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 2 ounces per 1000 cubic feet of space above the animals. Direct spray towards the upper portions of the enclosure filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment if reinfestation occurs.

KITCHEN PRODUCTION AND PROCESSING:
Control: Mosquitoes, Flies, Gnats
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply in kitchen houses at 1 ounce per 1000 cubic feet of space. Fly paper should be placed 8 to 10 feet away from beds and treated lightly upwards over them. Circulating fans should be left in operation. For best results, refrigerators should be turned off. In space sim. plants, use at rate of 1 ounce per 1000 cubic feet of space.

SPACE SPRAY APPLICATION:
Control: Mosquitoes, Flies, Gnats
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 2 ounces per 1000 cubic feet of space.

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

TREATMENT OF STORED GRAIN:
 To control a possible adult stages of:
 Indian meal moths, Angoumois grain beetle

FOR IN-TRANSIT PROTECTION: Clean and spray truck beds, boxcars, and ships' holds before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked.

TREATMENT OF STORED GRAIN:
 To control a possible adult stages of:
 Indian meal moths, Angoumois grain beetle

Apply to the surface of binned grain at a rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

TREATMENT OF HIDEAWAY INSECT NESTING PLACES:

To aid in the control of:
 Mosquitoes, Gnats, Wasps, Flippers
 Treat walls, ceiling, rafters, screens, door frames, window frames, beams, light fixtures and similar nesting places with the above dilution of M.A.G.C. 10-1.

CONTROL OF BEATING, UNCLE TOM:

Control: Mosquitoes, Small flying moths, Gnats

Apply to the surface of truck bed, equipment, and other areas as light breeze. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

SEE NEXT PAGE

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

CONTROL OF STORED GRAIN:
 Mix 14 ounces of Pyrethrin M.A.G.C. 10-1 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up the webbing with a rake before spraying and make a second application in 2 weeks.

CODE 720.05 PY NONE M.A.C.C. 10-1 CONTINUED

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

ENVIRONMENTAL PRECAUTIONS

This product is toxic to fish. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where there are important resources.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Do not pour or spill near open flame. Remove pets, birds and cover fish aquaria before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not apply directly to food. Do not apply space spray while food processing is underway. Foods should be removed or covered before treatment. Thoroughly wash all food processing surfaces before reuse. After spraying in meat packing plants, bakeries and food processing plants, wash with potable water, all equipment, benches, mixing, etc. where exposed food will be handled. If washing is impractical, do not allow food to come in contact with treated surfaces for at least 48 hours. Food processing operations do not have to be stopped while applying space spray with care and in accordance with the directions and cautions given above.

When using in animal quarters, do not apply directly to food, water or food supplements. Wash teats of dairy animals before milking.

Buyers should read directions of use, storage or handling of this product and if strict adherence to all directions given herewith.

BATCH

31

22