Reg # 11516-568 PM-17 131

CODE 720.05

Pyrenone®

M.A.G.C. 10-1

EPA Reg. No. 4816-568 EPA Est. No. 279-NY-1

For indoor or outdoor application as a space, area or contact spray. Dependent upon pests to be controlled and the area to be treated, may be applied through Mechanical Aerosol or ULV Generators, Thermal Fogging Equipment as well as other conventional fogging or spraying equipment. Designed for use in all areas of official establishments operating under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs.

INDOOR USE AREAS

Bakeries	Hotels	Storage or Processing Areas of Dried Food Products
Beverage Plants	Industrial Installations	Surfaces of Bags of Stored Food Products
Canneries	Meat Packing Plants	Supermarkets
Flour Mills	Motels	Truck Trailers
Food Processing Plants	Office Buildings	Warehouses
Grain Elevators	Railroad Cars	Ships' Holds
Granaries	Homes	Restaurants
Hospitals	Schools	

OUTDOOR USE AREAS

Cattle Barns	Golf Courses	Recreational Areas
Corrals	Horse Barns	Swine Houses
Drive-in Restaurants	Parks	Swine Lots
Drive-in Theaters	Playgrounds	Urban Areas
Feedlots	Poultry Houses	Zoos

ACTIVE INGREDIENTS

Pyrethrins	1.0%
*Piperonyl Butoxide Technical	10.0%
**INERT INGREDIENTS	89.0%
	100.0%

*Equivalent to min. 8% (butylcarbityl) (6-propylpiperonyl) ether and 2% related compounds

**Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE REAR PANEL FOR USE CAUTIONS

PYRENONE — Registered Trademark of Fairfield American Corporation.

NET CONTENTS: _____

BATCH

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Pyrenone M.A.G.C. 10-1 effectively controls the following insects when the applicator follows the directions for use as given below:

Angoumois Grain Moths	Drugstore Beetles	Mediterranean Flour Moths
Ants	Earwigs	Mushroom Flies
Boxelder Bugs	Fruit Flies	Red Flour Beetles
Cadelles	Fungus Gnats	Rice Weevils
Cheese Mites	Gnats	Sawtoothed Grain Beetles
Cheese Skippers	Grain Mites	Silverfish
Cigarette Beetles	Granary Weevils	Small Flying Moths
Clover Mites	Horn Flies	Spiders
Cockroaches	Hornets	Spider Beetles
Confused Flour Beetles	Horse Flies	Stable Flies
Crickets	House Flies	Tobacco Moths
Dark Mealworms	Indian Meal Moths	Wasps
Deer Flies	Lice	Yellow Mealworms
Dried Fruit Beetles	Mosquitoes	

ACCEPTED
APR 1 1988
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4816-568

INDOOR USE, UNDILUTED

This use of the product in food processing or food handling establishments (bakeries, beverage plants, canneries, food processing plants, hospitals, hotels, industrial installations, meat packing plants, restaurants, schools, supermarkets and warehouses) should be confined to time periods when the plants or facilities are not in operation. This use is not appropriate at any time in edible product areas of Federally inspected meat and poultry plants. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

SPACE AND CONTACT SPRAY APPLICATION

To control accessible adult stages of:

Flies	Hornets	Earwigs
Fruit Flies	Clover Mites	Ants
Mosquitoes	Cheese Mites	Cockroaches
Small Flying Moths	Cheese Skippers	Crickets
Wasps	Boxelder Bugs	Silverfish

and the following common stored product pests

Granary Weevils	Drugstore Beetles	Yellow Mealworms
Rice Weevils	Angoumois Grain Moths	Dark Mealworms
Confused Flour Beetles	Mediterranean Flour Moths	Grain Mites
Sawtoothed Grain Beetles	Indian Meal Moths	Red Flour Beetles
Spider Beetles	Tobacco Moths	Cadelles
Cigarette Beetles		

Close doors and windows and shut off ventilating systems. Use a mechanical aerosol generator or a fogger adjusted to deliver aerosol droplets (30 microns or less). Direct the spray first into all cracks and crevices, behind all equipment, cupboards, obstructions or dark harborage areas to provide maximum penetration and impingement on the insects. Then treat as a space spray, directing the spray towards the ceiling and upper corners of the area and behind any obstructions, apply at a rate of 0.9 oz. per 1000 cubic feet of space. Keep area closed for at least one hour. Vacate treated area and ventilate before reoccupying. Repeat treatment as necessary.

TREATMENT OF STORED FOOD AREAS

Foods in multi-walled bags or in cloth bags stored in warehouses, storage rooms, flour mills, grain elevators and granaries.

To control accessible adult stages of:

Angoumois Grain Moths	Sawtoothed Grain Beetles	Grain Mites
Cadelles	Yellow Mealworms	Granary Weevils
Confused Flour Beetles	Cheese Mites	Dark Mealworms
Indian Meal Moths	Cheese Skippers	Cockroaches
Mediterranean Flour Moths	Cigarette Beetles	Red Flour Beetles
Rice Weevils		

SEE REVERSE SIDE

FAIRFIELD AMERICAN CORPORATION
[illegible] WILSON AVENUE
NEWARK, N.J. 07105

RECEIVED
MAR 1988