

BEST AVAILABLE COPY

ACCEPTED
JUN 11 1985

**FAI
AM
CORP**
238 W
NEWA

SEE NEXT PAGE

OUTDOOR APPLICATION, DILUTED

TO CONTROL MOSQUITO LARVAE: Mix 18 ounces of Pyrenone M.A.G.C. 10-3.34 with sufficient oil to equal 50 gallons of diluted spray. Apply 20 to 25 gallons per acre. Apply as a coarse wetting spray to the border or on the shallow water of wet lands, swamps and marshes known to be infested with mosquito larvae. ANIMAL QUARTER USE (Cattle barns, horse barns, poultry houses, swine houses, zoos): As a space spray, diluted for use in conventional mechanical fogging or spraying equipment.

To control:
Flies
Small flying moths

Mix 6 ounces of Pyrenone M.A.G.C. 10-3.34 with sufficient oil to equal 1 gallon of diluted spray. Apply at the rate of 2 ounces per 1000 cubic feet of space above the animals. Direct spray at the upper portions of the enclosure filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before recouping. Repeat treatment if reinfestation occurs. In zoos, avoid exposure of reptiles to this product.

FOR USE ON ANIMALS

To protect against:

Horn flies	House flies
Mosquitoes	Onats

Mix 5 ounces of Pyrenone M.A.G.C. 10-3.34 with sufficient oil to equal 1 gallon of diluted spray. Apply a light mist sufficient to wet the tips of the hair on beef and dairy cattle, horses and dogs.

To protect against:

Stable flies	Horse flies
Deer flies	

Apply the above dilution at 2 ounces per adult beef cattle, dairy cattle and horses to sufficiently wet the hair, but not to soak the hide. Repeat this application once or twice daily or as needed to give continued protection.

To control the motile stages of blood sucking lice on beef and dairy cattle, apply the dilute spray to thoroughly wet the hair. Use two ounces or less per animal and repeat every 2 to 3 weeks as required.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spills or leakage, soak up with an absorbent material such as sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.
PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Remove pets, birds and caviar fish aquariums before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the United States Bureau of Mines to give adequate protection against this material. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not apply as a space spray while food processing is underway. Except in Federally inspected meat and poultry plants, when applied as a surface spray with care and in accordance with the directions and precautions given above food processing operations, may continue. Do not apply at any time as an undiluted space spray in edible product areas of Federally inspected meat and poultry plants. Foods should be removed or covered before treatments. All surfaces in processing areas must be washed and rinsed in potable water after spraying.

When using animal quarters, do not apply directly to food, water or food supplements. Wash teats of dairy animals before milking.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Contact a physician or Poison Control Center immediately. Do not induce vomiting unless directed to by a physician. Aspiration is a potential hazard.

If in Eyes: Flush with plenty of water. Contact a physician if irritation persists.

If on Skin: Wash affected areas with plenty of soap and water.

If Inhaled: Remove to fresh air. If breathing is difficult, seek medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, bays, marshes or estuaries. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

PHYSICAL OR CHEMICAL HAZARDS

Do not pour or spill near open flame.

Excess issues a risk of use, at rate or handling of this product not in strict accordance with directions given herein.

BATCH

Pyrenone®

M.A.G.C. 10-3.34

For trained personnel only

To be applied only by or under the supervision of Pest Control Operators, members of Mosquito Abatement Districts or Public Health Organizations and other trained personnel responsible for Insect Control Programs.

For indoor and outdoor application as a space, area or contact spray. Dependent upon pests to be controlled and the area to be treated, may be applied through mechanical aerosol or ULV Generators as well as conventional spraying (including thermal) or spraying equipment. Designed for use in all areas of official establishments operating under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs.

INDOOR USE AREAS:

Bakeries	Railroad Cars
Beverage Plants	Homes
Canneries	Schools
Flour Mills	Storage or Processing
Food Processing Plants	Areas of Dried
Grain Elevators	Food Products
Granaries	Surfaces of Bins of
Hospitals	Stored Food Products
Hotels	Supermarkets
Industrial Installations	Truck Trailers
Meat Packing Plants	Warehouses
Offices	Top of Halls
Office Buildings	Restaurants

OUTDOOR USE AREAS:

Cattle Bars	Playgrounds
Corrals	Poultry Houses
Drive-in-Restaurants	Recreational Areas
Drive-in-Theaters	Swine Houses
Feedlots	Swine Lots
Golf Courses	Urban Areas
Horse Barns	Zoos
Parks	

ACTIVE INGREDIENTS

Pyrethrins	3.34%
*Piperonyl Butoxide, Technical	10.00%
**INERT INGREDIENTS	86.66%
	100.00%

Equivalent to 3.0% (butylchrysinyl)X5-propyl-
pyrethrin ester and 2.0% related compounds

*Contains Petroleum Distillate
PYRENONE - Registered trademark of Fairfield
American Corporation

EPA Reg. No. 4419-571 EPA Est. No. 279-NY-1

NET CONTENTS

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See rear panel for additional precautions

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Pyrenone M.A.G.C. 10-3.34 effectively controls the following insects while the applicator follows the directions for use as given below:

Angoumois grain moth	Horn flies
Ants	Hornets
Boxelder bugs	Horse flies
Cadellies	House flies
Cheese mites	Indian meal moths
Cheese skippers	Lo
Cigarette beetles	Mosquitoes
Clover mites	Mediterranean flour moth
Cockroaches	Mushroom flies
Confined flour beetles	Red flour beetles
Crickets	Rice weevils
Dark mealworms	Sawtoothed grain beetles
Deer flies	Silverfish
Dried fruit beetles	Small flying moths
Drugstore beetle	Spiders
Earwigs	Spider beetles
Fruit flies	Stable flies
Fungus gnats	Tobacco moths
Gnats	Wasps
Grain mites	
Granary weevils	

INDOOR USE, UNDILUTED

This use of the product in food processing or food handling establishments (bakeries, beverage plants, canneries, food processing plants, hospitals, hotels, industrial installations, meat packing plants, restaurants, schools, supermarkets and warehouses) should be confined to time periods when the plants or facilities are not in operation. This use is not appropriate at any time in edible product areas of Federally inspected meat and poultry plants. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

SPACE AND CONTACT SPRAY APPLICATION

To control accessible adult stages of:

flies	Cheese skippers
fruit flies	Boxelder bugs
Mosquitoes	Earwigs
Small flying moths	Ants
Wasps	Cockroaches
Hornets	Crickets
Clover mites	Silverfish
Cheese mites	

and the following common stored product pests:

Granary weevils	Mediterranean flour
Rice weevils	moths
Confined flour beetles	Indian meal moths
Sawtoothed grain beetles	Tobacco moths
Spider beetles	Yellow mealworms
Cigarette beetles	Dark mealworms
Drugstore beetles	Grain mites
Angoumois grain moths	Red flour beetles
	Cadellies

Close doors and windows and shut off ventilating systems. Use a mechanical aerosol generator or a fogger adjusted to deliver aerosol droplets (30 microns or less). Direct the spray first into all cracks and crevices, behind all equipment, cupboard, obstructions or dark harborage areas to provide maximum penetration and impingement on the insects. Then treat as a space spray, directing the spray towards the ceiling and upper corners of the area and behind any obstructions, apply at a rate of 0.5 oz. per 1000 cubic feet of space. Keep area closed for at least one hour. Vacate treated area and ventilate before re-occupying. Repeat treatment as necessary.

TREATMENT OF STORED FOOD AREAS

Foods in multi-walled bags or in cloth bags stored in warehouses, storage rooms, flour mills, grain elevators and granaries.

To control accessible adult stages of:

Angoumois grain moths	Cheese mites
Cadellies	Cheese skippers
Confined flour beetles	Cigarette beetles
Indian meal moths	Grain mites
Mediterranean flour moth	Granary weevils
Rice weevils	Dark mealworms
Sawtoothed grain beetles	Cockroaches
Yellow mealworms	Red flour beetles

SEE REVERSE SIDE

ACCEPTED
JAN 31 1985
U.S. DEPT. OF AGRICULTURE
NATIONAL PEST CONTROL BOARD
WASHINGTON, D.C. 20250
FBI/DOJ
FBI/DOJ
FBI/DOJ

**FAIRFIELD
AMERICAN**

**FAIRFIELD
AMERICAN**

**FAIRFIELD
AMERICAN
CORPORATION**
238 WILSON AVENUE
NEWARK, N.J. 07102

BEST AVAILABLE COPY