

ALLEVIATE[®] FOGGING SPRAY

EPA Reg. No. 4816-579 EPA Est. No. 279-NY-1

ALLEVIATE - Registered Trademark of Fairfield American Corporation

For Insect Control In Food Plants, Restaurants, Hotels, Motels, Hospitals, Schools, Office Buildings, Warehouses, Boxcars, Ships' Holds, and Barns.

ACTIVE INGREDIENTS:

d/l-allethrolone d-trans chrysanthemate 0.50%
Related Compounds 0.04%

*Piperonyl Butoxide, Technical 5.00%

**INERT INGREDIENTS: 94.46%
100.00%

Equivalent to 4.0% (butylcarbityl) (6-propyl-piperonyl) ether and to 1.0% related compounds.

**Contains Petroleum Distillate.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See side panel for additional precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply directly to food. Do not apply space spray while food processing is underway. Foods should be removed or covered before treatment. Thoroughly wash all food processing surfaces before reuse. After spraying in meat processing plants, bakeries, and the food processing plants, wash with potable water all equipment, benches, shelving, etc. where exposed food will be handled. In case washing is impractical, do not allow food to come in contact with treated surfaces for at least 48 hours. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions given above.

For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths.

Close windows and doors and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. In Official Meat and Poultry Establishments use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. In other than Official

Meat and Poultry Establishments a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites, and cadellies: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs and infested clothing and furniture. Make a localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as necessary.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storage containers, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly, keeping the spray nozzle at least three feet from the fabric to avoid staining. Store clothes in moth tight containers or repeat treatments monthly. In Horse Barns and on Horses: This insecticide may be applied as a fine mist spray to stable area and over and around stabled horses for fast paralysis and kill of flies. Do not wet horse's skin or exceed two ounces per application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by an approved waste disposal facility.

For Containers One Gallon And Smaller

STORAGE: Store in original container in a cool dry place inaccessible to children and pets.

DISPOSAL: Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

BEST AVAILABLE COPY

CONTINUED ON REVERSE

Not Reviewed. Registrant claims to be in accordance with
FEDERAL REGISTER, Vol. 15, No. 100, for Phase 2
of the Pesticide Labeling Act, 1972, in Status

**FAIRFIELD
AMERICAN
CORPORATION**

201 Route 17 North, Rutherford, N.J. 07070